# 

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS14 2774 Anti-ACC synthase 7 | 1-aminocyclopropane-1-carboxylate synthase 7


### **Product information**

Immunogen	KLH-conjugated synthetic peptide derived from Arabidopsis thaliana ACC synthase 7, UniProt: Q9STR4,TAIR: AT4G26200	
Host	Rabbit	
Clonality	Polyclonal	
Purity	Immunogen affinity purified serum in PBS pH 7.4.	
Format	Lyophilized	
Quantity	100 µg	
Reconstitution	For reconstitution add 100 $\mu$ l of sterile water	
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.	
Additional information	ACS7 protein levels are too low to allow detection in endogenous cell extract	
Application information		

#### **Application information**

Recommended dilution	1 : 500 (WB)
Expected   apparent MW	50.6 kDa
Confirmed reactivity	Arabidopsis thaliana
Predicted reactivity	Arabidopsis thaliana
Not reactive in	No confirmed exceptions from predicted reactivity are currently known

#### Application example



10 ug of total protein extract from *Nicotiana benthamiana* leaves expressed ACS7–His–GST (1), 5 ug of purified ACS7–His–GST from *N. benthamiana* ACS7–His–GFP fusion protein transiently expressed in *N. benthamiana* leaves was purified on Ni NTA resin(2) and (3). Samples were separated on 10% SDS–PAGE and transfered to PVDP membrane by semi–dry blotting (1h). Blot was blocked with 3% skim milk in PBS–T (0,01% Tween 20) for 1h in RT with agitation. Then blot was incubated with primary antibody anti ACS7 at dilution of 1:500 in PBS–T for 1h in RT with agitation. After washes (3 times for 5 min. in PBS–T) blot was incubated with secondary antibody (anti–rabbit IgG horse radish peroxidase conjugated, from Agrisera, <u>AS09 602</u>) at dilution 1:25000 for 1h in RT. The blot was washed 5 times for 5 min. with PBS–T and developed for 2 min with ECL–Plus according to the manufacturer's instructions. Exposure time was 3 min.

Courtesy of Dr. Agata Cieśla, Ludwików's lab, UAM, Poland