


Product no **AS10 932****Anti-Amyloid beta oligomer-specific monoclonal antibody (OMAB)****Product information**


Immunogen	Partly aggregated, recombinant peptide corresponding to the human Abeta (1-40/42). Amino acid sequence: D-A-E-F-R-H-D-S-G-Y-E-V-H-H-Q-K-L-V-F-F-A-E-D-V-G-S-N-K-G-A-I-I-G-L-M-V-G-G-V-V. The epitope is 3-8. The molecular weight of immunogen is 4.5 kDa.
Host	Mouse
Clonality	Monoclonal
Subclass/isotype	IgM
Purity	Affinity purified in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 100 µl of sterile water
Storage	Store lyophilized/reconstituted at 4 °C. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
Additional information	OMAB antibody has been purified by ion-exchange chromatography and is supplied in PBS without any additives as carrier proteins or sodium azide. Binding of OMAB antibody and Abeta oligomers at RT takes about 15 min. Fibrils are inaccessible for OMAB antibodies therefore if a discrimination between fibrils and oligomers is to be achieved, dot blot can be used. Start with antigen concentration of 500 ng/dot followed by 2X dilution steps. Blocking: non-fat milk and washes with 0.3 % Tween 20 in TBS pH 7.4.

Application information

Recommended dilution	Coating antibody at 2 µg/ml (ELISA), 1: 1000 (IF), 1 : 500 (IHC)
Expected apparent MW	4.5 kDa
Confirmed reactivity	Human Abeta oligomers only
Predicted reactivity	Rat
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	OMAB antibody is a versatile tool within research of Alzheimer's disease, A sandwich ELISA illustrates its potential regarding its high selectivity towards A β oligomers
Selected references	Nagashima et al. (2024). Development of Triphenylmethane Dyes for In Vivo Fluorescence Imaging of A β Oligomers. ACS Chem Neurosci. 2024 Jun 5;15(11):2233-2242. doi: 10.1021/acscchemneuro.4c00053. Akasanka et al. (2023). In Vivo Near-Infrared Fluorescence Imaging Selective for Soluble Amyloid Aggregates Using γ -Shaped BODIPY Derivative. J Med Chem. 2023 Oct 26;66(20):14029-14046. doi: 10.1021/acs.jmedchem.3c01057. Pang et al (2021) An App knock-in rat model for Alzheimer's disease exhibiting A β and tau pathologies, neuronal death and cognitive impairments. Cell Res. 2021 Nov 17. doi: 10.1038/s41422-021-00582-x. Epub ahead of print. PMID: 34789895. Oh et al. (2020). Associative Interactions among Zinc, Apolipoprotein E, and Amyloid- β in the Amyloid Pathology. Int J Mol Sci. 2020 Jan 25;21(3). pii: E802. doi: 10.3390/ijms21030802. Henning-Knechtel et al. (2020). Designed Cell-Penetrating Peptide Inhibitors of Amyloid-beta Aggregation and Cytotoxicity. Cell Reports Physical Science, Volume 1, Issue 2, 26 Zhang et al. (2019). Brains of rhesus monkeys display A β deposits and glial pathology while lacking A β dimers and other Alzheimer's pathologies. Aging Cell. 2019 Jun 4:e12978. doi: 10.1111/acer.12978. Kumar et al. (2018). Peptidomimetic-Based Multidomain Targeting Offers Critical Evaluation of A β Structure and Toxic Function. J Am Chem Soc. 2018 May 30;140(21):6562-6574. doi: 10.1021/jacs.7b13401.



Abeta oligomer-specific antibody was adsorbed to Nunc-Immuno MaxiSorp plates (Nunc, Roskilde, Denmark) at 2 ug/ml in PBS. 1 ml of a 10 uM A β (1-42) sample containing a small fraction of A β -oligomers was separated using a superdex G75 (10/30) column. A β -fractions collected from the SEC were allowed to bind to OMAB plates for 20 minutes at 0°C. All fractions were analyzed and bound A β was detected using a polyclonal rabbit anti-A β antibody ([AS08 328](#), Agrisera AB, Vännäs, Sweden) at a 1:1000 dilution followed by an anti-rabbit HRP-conjugated secondary antibody at a 1:5000 dilution (GE healthcare). ECBlue (Medicago, Uppsala, Sweden) was used as a substrate for HRP and the signal was detected by measuring the absorbance at 450 nm. Blocking solution and antibody-dilutions were made with 5% Non-fat dry milk in PBST and all washes were performed with PBS containing 0.1% Tween-20 (PBST).



10 μ m of coronal sections from fresh-frozen transgenic mouse brain mutant (**A**) and wild type (**B**). Post-fixation in 4% formaldehyde solution, 5 min. OMAB antibody diluted 1:500, incubation at 4°C ON. Mouse on mouse HRP-Polymer kit according to company instructions. Biocare Medical: BC-MM510 (Histolab) DAB substrate kit for peroxidase. Vector Laboratories: SK-4100 (ImmunKemi) Counterstained with Mayers HTX.