


Product no **AS11 1810****Anti-RACK1A | Receptor for activated C kinase 1A****Product information**

| | |
|-----------------------|---|
| Immunogen | KLH-conjugated synthetic peptide derived from <i>Arabidopsis thaliana</i> RACK1A protein sequence UniProt: O24456 , TAIR: At1g18080 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Purity | Serum |
| Format | Lyophilized |
| Quantity | 50 µl |
| Reconstitution | For reconstitution add 50 µl of sterile water |
| Storage | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube. |

Application information

| | |
|-------------------------------|--|
| Recommended dilution | 1 : 2000 (WB) |
| Expected apparent MW | 35 kDa |
| Confirmed reactivity | <i>Arabidopsis thaliana</i> |
| Predicted reactivity | <i>Thellungiella salsuginea</i> |
| | Species of your interest not listed? Contact us |
| Not reactive in | <i>Nicotiana benthamiana</i> , <i>Zea mays</i> |
| Selected references | Hemayet et al. (2019) . Host targeted antiviral (HTA): functional inhibitor compounds of scaffold protein RACK1 inhibit herpes simplex virus proliferation. <i>Oncotarget</i> . 2019 May 14; 10(35): 3209–3226. Vera-Estrella et al. (2014) . Comparative 2D-DIGE analysis of salinity responsive microsomal proteins from leaves of salt-sensitive <i>Arabidopsis thaliana</i> and salt-tolerant <i>Thellungiella salsuginea</i> . <i>J Proteomics</i> . 2014 Jun 2. pii: S1874-3919(14)00288-7. doi: 10.1016/j.jprot.2014.05.018. Speth et al. (2013) . RACK1 scaffold proteins influence miRNA abundance in <i>Arabidopsis</i> . <i>Plant J</i> . Aug 13. |

application example

20 µg of total protein from *Arabidopsis thaliana* total cell extract was separated on **15 % SDS-PAGE** and blotted 1h to **PVDF**. Blots were blocked with Roti-block over night at 4°C agitation. Blot was incubated in the primary antibody at a dilution of 1: 1 000 for 1h at RT with agitation. Blot was incubated in the primary antibody at a dilution of 1: 2 000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera [AS09 602](#)) diluted to 1:10 000 in for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL according to the manufacturers instructions. Exposure time was 30 min.

Courtesy of Dr. Sascha Laubinger, ZMBP, Germany