

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS11 1737

## Anti-Beta-CA1, beta-CA2 | carbonic anhydrase | mitochondrial | Chlamydomonas

## **Product information**

Immunogen recombinant Chlamydomonas reinhardtii mitochondrial CA, as described in Villand et al. 1997. Accession number Q39590 and Q39589

**Host** Rabbit

Clonality Polyclonal

**Purity** Serum

Format Lyophilized

Quantity 200 ul

**Reconstitution** For reconstitution add 200 μl of sterile water

Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please Storage remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to

the cap or sides of the tube.

Antibody is recognizing both isoforms, beta- CA1 and beta-CA2 and can be used as mitochondrial marker for low Additional information carbon dioxide grown cells of Chlamydomonas reinhardtii

## Application information

Recommended dilution 1:200 (IF), 1:1000 (WB)

Expected | apparent

23.7 | 21-22 kDa

Confirmed reactivity Chlamydomonas reinhardtii

Predicted reactivity Chlamydomonas reinhardtii

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Selected references

Burlacot et al. (2022) Alternative photosynthesis pathways drive the algal CO2-concentrating mechanism. Nature 605, 366-371 (2022). https://doi.org/10.1038/s41586-022-04662-9

Kuken et al. (2018). Effects of microcompartmentation on flux distribution and metabolic pools in Chlamydomonas reinhardtii chloroplasts. Elife. 2018 Oct 11;7. pii: e37960. doi: 10.7554/eLife.37960.

Muranaka et al. (2015). TEF30 interacts with photosystem II monomers and is involved in the repair of photodamaged photosystem II in Chlamydomonas reinhardtii. Plant Physiol. 2015 Dec 7. pii: pp.01458.2015.

Tirumani et al. (2014). Regulation of CCM genes in Chlamydomonas reinhardtii during conditions of light-dark cycles in synchronous cultures. Plant Mol Biol. 2014 Mar 4.

Renberg et al. (2010). A Metabolomic Approach to Study Major Metabolite Changes during Acclimation to Limiting CO2 in Chlamydomonas reinhardtii. Plant physiol. 154: 187-196.