


Product no **AS13 2641****Anti-CGL40****Product information**

Immunogen	Full-length, recombinant CGL40, UniProt: Q8LFL5 , TAIR: At1g49975
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	14.73 kDa-cTP: 8.42 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Capsella rubella</i> , <i>Brassica pekinensis</i>
	Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known

Application example

3 µg of total protein from *Arabidopsis thaliana* extracted with were separated on 15% SDS-PAGE and blotted 1h to PVDF. Blots were blocked with 10% skimmed milk for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1 000 overnight at 4°C with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed twice for 15 min and 4 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (goat anti-rabbit IgG, HRP-conjugated) diluted to 1:15 000 in TBS-T for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL according to the manufacturer's instructions. Exposure time was 30 seconds.

Antibody incubation can be also conducted for 1 h/RT.

Courtesy of Dr. Rikard Fristedt, UCLA, USA