

Product no **AS10 1112**

Chicken anti-Goat IgG (H&L), ALP conjugated

Product information

Host | Chicken**Clonality** | Polyclonal**Purity** | Immunogen affinity purified IgY.**Format** | Liquid**Quantity** | 1 mg**Storage** | Non-diluted antibody is stable for 4 years at 2-8°C. For storage at -20°C dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50 % to prevent loss of enzymatic activity. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.**Additional information** | This antibody reacts with the heavy chains on goat IgG and with the light chains on all goat immunoglobulins based on immunoelectrophoresis.

Minimum cross-reactivity is observed to non-immunoglobulin goat serum proteins.

Antibody is supplied in 30 mM triethanolamine, pH 7.2, 5 mM magnesium chloride, 0.1 mM zinc chloride, 1 % (w/v) BSA, Protease/IgG free, 0.05 % (w/v) sodium azide as preservative.

BSA and milk have to be replaced by other blocking reagents, like donkey serum or commercial formulations which are free from bovine IgG.

Application information

Recommended dilution | 1 : 500-1 : 2000 (ELISA), 1 : 50-1 : 5000 (ICC), 1 : 20-1 : 2000 (IHC), 1 : 500-1 : 2000 (WB)