Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com


Product no AS09 601 Anti-c-Myc (polyclonal)

Product information

Immunogen	KLH-conjugated peptide, sequence MEQKLISEEDLNE, human c-Myc UniProt: Q6LBK7
Host	Chicken
Clonality	Polyclonal
Purity	Immunogen affinity purified IgY in PBS pH 8.5. Contains 0.02% azide.
Format	Lyophilized
Quantity	100 μg
Storage	Store lyophilized antibodies at -20°C and reconstituted antibodies at 4°C. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
	-

Application information

Recommended dilution	1 : 2000-1 : 5000 (ICC)
Expected apparent MW	Depends on target protein which is tagged with this epitope
Confirmed reactivity	с-Мус
Predicted reactivity	с-Мус
Not reactive in	No confirmed exceptions from predicted reactivity are currently known



HEK293 cells were transfected with the indicated plasmid (SARS-CoV-2 N myc tagged, PMID: 34799561, TBEV NS3-HA, PMID: 29321318, Rab5 mcherry, Addgene: 4920, GFP-HIS) using genejuice transfection reagent (EMD Millipore) according to the manufacturer's instructions. After 24 hours of transfection, cells were fixed in 4% formaldehyde and permeabilized in PBS containing 0.5% Triton X-100 and 20 mM glycine. Then, cells were stained with the primary anti-Myc tag antibodies at a concentration of 1 µg/mL for 1 hour at room temperature. Followed by three washes in PBS. Cells were then stained using secondary antibodies, goat anti-chicken Alexa488 (1:500, Invitrogen A11039) in PBS containing 2% BSA for 1h at RT. Nuclei were stained using DAPI (1 µg/mL). Images were acquired using a Leica SP8 Laser Scanning Confocal Microscope with a 63x oil objective (Leica) and Leica Application Suit X software (LAS X, Leica).

Courtesy of Dr. Anna K Överby, Molecular Infection Medicine Sweden (MIMS), Section of Virology. Department of Clinical Microbiology Umeå University, Sweden

contact: support@agrisera.com