

Product no **AS10 111S****CP43' | IsiA homolog of plant CP43 positive control/quantitation standard****Product information****Format** | Lyophilized**Quantity** | 250 µl**Reconstitution** | For reconstitution add 225 µl of milliQ water**Storage** | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.**Additional information** | The IsiA protein standard can be used in combination with anti-IsiA antibodies to quantitate IsiA from a range of cyanobacteria. [Global antibodies](#) are raised against highly conserved amino acid sequences in the IsiA protein.Quantitative western blot: [detailed method description](#), [video tutorial](#)**Application information****Recommended dilution** | Standard curve: 3 loads are recommended (2.5 and 10 µl).  
For most applications a sample load of 0.2 µg of chlorophyll will give a IsiA signal in this range.

Positive control: a 2 µl load per well is optimal for most chemiluminescent detection systems.

This standard **is stabilized and ready** and does not require heating before loading on the gel. Please note that this product contains 10% glycerol and might appear as liquid but is provided lyophilized. Allow the product several minutes to solubilize after adding water. Mix thoroughly but gently. Take extra care to mix thoroughly before each use, as the proteins tend to settle with the more dense layer after freezing.**Expected | apparent MW** | 27 kDa (slightly larger than native protein due to His-tag)**Additional information** | **Concentration:** after adding 225 µl of milliQ water final concentration of the standard is 0.15 pmoles/µl**Protein standard buffer composition:** Glycerol 10%, Tris Base 141 mM, Tris HCl 106 mM, LDS 2%, EDTA 0.51 mM, SERVA® Blue G250 0.22 mM, Phenol Red 0.175 mM, pH 8.5, 0.1 mg/ml PefaBloc protease inhibitor (Roche), 50mM DTT.**This standard is ready-to-load and does not require any additions or heating. It needs to be fully thawed and thoroughly mixed prior to using. Avoid vigorous vortexing, as buffers contain detergent. Following mixing, briefly pulse in a microcentrifuge to collect material from cap.****Please note that this product contains 10% glycerol and might appear as liquid but is provided lyophilized. Allow the product several minutes to solubilize after adding water. Mix thoroughly but gently. Take extra care to mix thoroughly before each use, as the proteins tend to settle with the more dense layer after freezing.****Selected references** | [Fraser et al. \(2013\)](#). Photophysiological and Photosynthetic Complex Changes during Iron Starvation in Synechocystis sp. PCC 6803 and Synechococcus elongatus PCC 7942. PLoS ONE 8(3): e59861. doi:10.1371/journal.pone.0059861  
[Ryan-Keogh et al. \(2012\)](#). Iron deficiency in cyanobacteria causes monomerization of photosystem I trimers and reduces the capacity for state transitions and the effective absorption cross section of photosystem I in vivo. J. of Phycology, 1:145-154.