

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS06 123 CPX1 | coproporphyrinogen III oxidase, isoform 1

Product information

Immunogenresidues 32-366 from mature coproporphyrinogen III oxidase, isoform CPX1 of Chlamydomonas reinhardtii fused to
TrxA Q9S7V1HostRabbitClonalityPolyclonalPuritySerumFormatLyophilizedQuantity100 μlReconstitutionFor reconstitution add 100 μl of sterile waterStorageStore lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please
remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to
the cap or sides of the tube.

Application information

••	
Recommended dilution	1 : 3000 (WB)
Expected apparent MW	41,4 38 kDa
Confirmed reactivity	Physcomitrium patens
Predicted reactivity	<i>Arabidopsis thaliana, Zea mays</i> Species of your interest not listed? <u>Contact us</u>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Selected references	Lang et al. (2011).Simultaneous isolation of pure and intact chloroplasts and mitochondria from moss as the basis for sub-cellular proteomics. Plant Cell Rep. Feb;30(2):205-15. (reactivity confirmed for Physcomitrella patens). Quinn et al. (1999) Induction of Coproporphyrinogen Oxidase in Chlamydomonas Chloroplasts Occurs via Transcriptional Regulation of Cpx1 Mediated by Copper-Response Elements and Increased Translation from a Copper-Deficiency-Specific Form of the Transcript. J. Biol. Chem. 274:14444-14454.