

Product no **AS09 437****Anti-Cytokinin | Dihydrozeatin riboside (for immunolocalization)****Product information**

Immunogen	BSA-conjugated, via ribose, dihydrozeatin riboside
Host	Rabbit
Clonality	Polyclonal
Purity	Total IgG. Protein G purified in PBS.
Format	Liquid
Quantity	200 µg
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Additional information | Cytokinin | Dihydrozeatin riboside (for immunolocalization)**Application information**

Recommended dilution	Specific information about dilution is going to be included on the vial
Confirmed reactivity	Dihydrozeatin riboside
Predicted reactivity	Dihydrozeatin riboside
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	For 1 mg of antibodies For ELISA and immunoassay or For ELISA kit - please inquire, Antibodies are supplied in 50% glycerol
Selected references	Alvarez et al. (2020) . Hormonal and gene dynamics in de novo shoot meristem formation during adventitious caulogenesis in cotyledons of Pinus pinea. Plant Cell Rep. 2020 Jan 28. doi: 10.1007/s00299-020-02508-0.