

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91112 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

## Product no AS07 206

### Anti-Dehydrin (serum)

#### Product information

<b>Immunogen</b>	KLH-conjugated peptide sequence TGEKKGIMDKIKEKLPGQH of K-segment conserved in a wide range of plant species
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µl
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile, deionized water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

#### Application information

<b>Recommended dilution</b>	1 : 1000 (WB)
<b>Expected   apparent MW</b>	9-200 kDa
<b>Confirmed reactivity</b>	<i>Agrostis stolonifera</i> cv. 'Penncross', <i>Betula pubescens</i> , <i>Betula pendula</i> , <i>Betula pendula</i> var. <i>carellica</i> , <i>Hordeum spontaneum</i> , <i>Larix cajanderi</i> , <i>Malus</i> spp., <i>Picea obovata</i> , <i>Picea abies</i> , <i>Picea glauca</i> , <i>Pinus sylvestris</i> , <i>Pinus strobus</i> , <i>Pinus sylvestris</i> , <i>Spinacia oleracea</i> , <i>Triticum aestivum</i> , <i>Vitis vinifera</i>
<b>Predicted reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Glycine max</i> , <i>Nicotiana tabacum</i> , <i>Pisum sativum</i> , <i>Hordeum vulgare</i> , <i>Oryza sativa</i> , <i>Populus</i> sp., <i>Zea mays</i>
	Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Additional information</b>	According to <a href="#">Borovskii</a> et al. 2019, dehydrin detection level can be increased by obtaining a thermostable fraction.
<b>Selected references</b>	<p><a href="#">Vitámvás</a> et al. (2021) Relationship between WCS120 Protein Family Accumulation and Frost Tolerance in Wheat Cultivars Grown under Different Temperature Treatments. <i>Plants</i> (Basel). 2021 May 31;10(6):1114. doi: 10.3390/plants10061114. PMID: 34073120; PMCID: PMC8228299.</p> <p><a href="#">Kartashov</a> et al. (2021) Quantitative analysis of differential dehydrin regulation in pine and spruce seedlings under water deficit. <i>Plant Physiol Biochem</i>. 2021 Mar 3;162:237-246. doi: 10.1016/j.plaphy.2021.02.040. Epub ahead of print. PMID: 33706184.</p> <p><a href="#">Vazquez-Hernandez</a> et al. (2020). Functional characterization of VviDHN2 and VviDHN4 dehydrin isoforms from <i>Vitis vinifera</i> (L.): An in silico and in vitro approach. <i>Plant Physiol Biochem</i>. 2021 Jan;158:146-157. doi: 10.1016/j.plaphy.2020.12.003. Epub 2020 Dec 4. PMID: 33310482.</p> <p><a href="#">Rachenko</a> and Rachenko (2020). The variation of the content of dehydrin proteins in the bark of <i>Malus</i> spp. trees differing in winter hardiness in Southern Cisbaikalia conditions. <i>Zemdirbyste-Agriculture</i>, vol. 107, No. 2 (2020), p. 185–190 DOI 10.13080/z-a.2020.107.024.</p> <p><a href="#">Lv</a> et al. (2018). Characterization of Dehydrin protein, CdDHN4-L and CdDHN4-S, and their differential protective roles against abiotic stress in vitro. <i>BMC Plant Biol</i>. 2018 Nov 26;18(1):299. doi: 10.1186/s12870-018-1511-2.</p>