

Product no **AS16 3244****Goat anti-Bovine IgG (H&L), Unconjugated****Product information**

Immunogen	Purified Bovine IgG, whole molecule
Host	Goat
Clonality	Polyclonal
Purity	Immunogen affinity purified in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.05% (w/v) sodium azide.
Format	Liquid
Quantity	2,0 mg
Storage	Store at 2-8°C.

Additional information	<p>Antibody purity is > 95% based on SDS-PAGE</p> <p>Affinity purified using solid phase Bovine IgG (H&L)</p> <p>Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2.</p> <p>0.05% (w/v) Sodium Azide is added as preservative.</p> <p>Based on IEP, this antibody reacts with: · heavy () chains on bovine IgG · light chains on all bovine immunoglobulins.</p> <p>Based on IEP, no reactivity is observed to: · non-immunoglobulin bovine serum immunoglobulins.</p>
-------------------------------	--

Application information

Recommended dilution	The optimal working dilution should be determined by the investigator
Confirmed reactivity	Bovine IgG (H&L)
Selected references	<u>Patathil</u> et al. (2010). A comprehensive toolkit of plant cell wall glycan-directed monoclonal antibodies. Plant Physiol. 2010 Jun;153(2):514-25.doi: 10.1104/pp.109.151985.