

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS16 3264

Goat anti-Chicken IgY (H&L), DyLight® 405 conjugated

Product information

Immunogen Purified Chicken IgY, whole molecule

Host Goat

Clonality Polyclonal

Purity Immunogen affinity purified goat IgG.

Format Lyophilized

Quantity 1 mg

Reconstitution

For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily

Storage

Store freeze-dried powder at 2-8 °C. Product is stable for 4 weeks at 2-8 °C after rehydration. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20 °C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20 °C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.

Additional information

Antibody purity is > 95% based on SDS-PAGE.

DyLight® 405 (Ex = 400 nm; Em = 420 nm) Affinity purified using solid phase Chicken lgG/Y.

Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/lgG free. 0.05% (w/v) Sodium Azide is added as a preservative.

Based on IEP, this antibody reacts with: • heavy () chains on chicken IgG/Y • light chains on all chicken immunoglobulins .

Based on IEP, no reactivity is observed to: • non-immunoglobulin chicken serum immunoglobulins .

Application information

Recommended dilution 1:20-1:2000 (Flow cyt), (IL)

Confirmed reactivity Chicken IgY

Selected references

Huiming et al. (2020). Integration of Nodal and BMP Signaling by Mutual Signaling Effector Antagonism. Cell Rep. 2020 Apr 7;31(1):107487.doi: 10.1016/j.celrep.2020.03.051.