

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS16 3123

Goat anti-Dog IgG Fc specific, HRP conjugated

Product information

Immunogen Purified normal dog IgG, isolated from pooled dog serum

Host Goat

Clonality Polyclonal

Purity Protein G purified goat IgG in PBS pH 7.2.

Format Lyophilized

Quantity 10 mg

Reconstitution For reconstitution add 1 ml of sterile destilled water

Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to

the cap or sides of the tube. Reconstituted material can be stoored in 4°C up to 7 days.

Application information

Recommended dilution 1:2000-1:10 000 (ELISA), 1:1001:-500 (ICC), (IH), The optimum working dilution should be established by titration by end user, Excess labelled antibody should be avoided since it may cause high unspecific background staining and interfere with the specific signal

Confirmed reactivity Dog IgG, Fc part

Not reactive in

This antibody does not react with any non-lg protein in dog serum, as tested by immunoelectrophoresis and double radial immunodiffusion

Additional information

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antibody has not been tested in detail.

No preservative added.