Agrisera Part of Olink® Group \$

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS16 3130 Goat anti-Dog IgM Fc specific, FITC conjugated

Product information

Immunogen	Purified normal IgM isolated from pooled dog serum
Host	Goat
Clonality	Polyclonal
Purity	Protein G purified goat IgG in PBS pH 7.2.
Format	Lyophilized
Quantity	10 mg
Reconstitution	For reconstitution add 1 ml of sterile destilled water
-	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube. Reconstituted material can be stoored in 4°C up to 7 days.

Application information

Recommended dilution	1 : 10-1 :40 (ICC), (IHC), (IF), The optimum working dilution should be established by titration by end user, Excess labelled antibody should be avoided since it may cause high unspecific background staining and interfere with the specific signal
Confirmed reactivity	dod IgM, Fc part
Not reactive in	This antibody does not react with IgG, IgG/Fab fragments and IgA or any non-Ig protein in dog serum, as tested by immunoelectrophoresis and double radial immunodiffusion
Additional information	This immunoconjugate is not species-specific since inter-species cross-reactivity is a normal feature of antibody to immunoglobulins. Cross-reactivity of this antibody has not been tested in detail. No preservative added.