

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS16 3130

## Goat anti-Dog IgM Fc specific, FITC conjugated

## **Product information**

Immunogen Purified normal IgM isolated from pooled dog serum

Host Goat

Clonality Polyclonal

**Purity** Protein G purified goat IgG in PBS pH 7.2.

Format Lyophilized

Quantity 10 mg

**Reconstitution** For reconstitution add 1 ml of sterile destilled water

Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube. Reconstituted material can be stoored in 4°C up to 7 days.

## **Application information**

Recommended dilution

1:10-1:40 (ICC), (IHC), (IF), The optimum working dilution should be established by titration by end user, Excess labelled antibody should be avoided since it may cause high unspecific background staining and interfere with the specific signal

Confirmed reactivity dod IgM, Fc part

Not reactive in

This antibody does not react with IgG, IgG/Fab fragments and IgA or any non-Ig protein in dog serum, as tested by immunoelectrophoresis and double radial immunodiffusion

**Additional information** 

This immunoconjugate is not species-specific since inter-species cross-reactivity is a normal feature of antibody to immunoglobulins. Cross-reactivity of this antibody has not been tested in detail.

No preservative added.