

Product no **AS12 1899****Goat anti-Human IgA (alpha chain), DyLight® 550 conjugated, min cross reactivity to bovine, mouse, rabbit serum****Product information**

| | |
|-------------------------------|---|
| Immunogen | purified human IgA (alpha chain) |
| Host | Goat |
| Clonality | Polyclonal |
| Purity | Immunogen affinity purified goat IgG. |
| Format | Lyophilized |
| Quantity | 1 mg |
| Reconstitution | For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily |
| Storage | Store lyophilized material at 2-8°C. Product is stable for 4 weeks at 2-8°C after rehydration. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1,1 ml of sterile water add 1,1 ml of glycerol. Such solution will not freeze in -20°C, If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming. |
| Additional information | Conjugate is present in 10 mM Sodium Phosphate, 0,15 M Sodium Chloride, pH 7,2, 1 % (w/v) BSA, Protease/IgG free, 0,05 % (w/v) sodium azide is added as preservative |

Application information

| | |
|-------------------------------|---|
| Recommended dilution | 1 : 20-1 : 2000 for most applications |
| Additional information | Based on immunoelectrophoresis, this antibody reacts with: heavy () chains on human IgA (chain) No reactivity is observed to: non-immunoglobulin human serum proteins light chains on all human immunoglobulins serum proteins from bovine, mouse or rabbit. |