

This product is for research use only (not for diagnostic or therapeutic use)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS16 3468 Goat anti-Mouse IgG (H&L), F(ab)'2 fragment, TRITC conjugated, min, cross-reactivity to bovine, goat, human, rabbit, or rat IgG

Product information

Immunogen	Purified Mouse IgG, whole molecule
Host	Goat
Clonality	Polyclonal
Purity	Immunogen affinity purified goat IgG.
Format	Lyophilized
Quantity	0,5 mg
	For reconstitution add 0,55 ml of sterile water,Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily
Storage	Store at 2-8°C.
Additional information	Antibody purity is > 90% based on SDS-PAGE Small amounts of intact IgG may be present. Affinity purified using solid phase Mouse IgG . Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. Based on IEP, this antibody reacts with: • heavy () chains on mouse IgG • light chains on all mouse immunoglobulins . Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • bovine, goat, human, rabbit or rat IgG .

Application information

Recommended dilution 1 : 20-1 : 2000 (Flow cyt), (IL) Confirmed reactivity Mouse IgG (H&L), F(ab)'2 Fragment