

Product no **AS10 969****Goat anti-mouse IgM (μ chain), HRP conjugated, min, cross-reactivity to human IgG/serum****Product information**

Host	Goat
Clonality	Polyclonal
Purity	Immunogen affinity purified goat IgG.
Format	Lyophilized
Quantity	1 mg
Reconstitution	For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily
Storage	Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1,1 ml of sterile water add 1,1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
Additional information	<p>This antibody reacts with the heavy chains on mouse IgM based on immunoelectrophoresis.</p> <p>Minimum cross-reactivity is observed with the light chains based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin mouse serum proteins or to human IgG or serum proteins based on immunoelectrophoresis.</p> <p>Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free and 0.1 % (v/v) Kathon CG is used as preservative. Use of sodium azide will inhibit enzymatic activity of horseradish peroxidase.</p>

Application information**Recommended dilution** 1 : 500-1 : 5000 (IHC), 1 : 10 000-1 : 50 000 (WB)