

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS11 1812

Goat anti-Rabbit IgG (H&L), DyLight® 350 conjugated

Product information

Immunogen Purified Rabbit IgG, whole molecule

Host Goat

Clonality Polyclonal

Purity Immunogen affinity purified goat IgG.

Format Lyophilized

Quantity 1 mg

Reconstitution

For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily

Storage

Store lyophilized material at 2-8°C. Product is stable for 4 weeks at 2-8°C after rehydration. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1,1 ml of sterile water add 1,1 ml of glycerol. Such solution will not freeze in -20°C, If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.

Additional information

Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative.

DyLight® 350 has a maximum absorbance at 353 nm; Emax = 432 nm.

Application information

Recommended dilution 1:50-1:5 000 (ICC), 1:20-1:2000 (IHC), 1:3000 (IF)

Confirmed reactivity Rabbit IgG heavy and light chains (H&L)

Predicted reactivity Rabbit IgG Heavy and Light chains (H&L)

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Additional information

Based in immunoelectrophoresis, this antibody reacts with heavy chains on rabbit IgG and light chains on all rabbit immunoglobulins.

No reactivity is observed to non-immunoglobulin rabbit serum proteins based in immunoelectrophoresis.

Purity of this antibody is > 95% based on SDS-PAGE.