

Product no **AS10 718****Anti-HTA9 | Probable histone H2A variant 3****Product information**

<b>Immunogen</b>	KLH-conjugated synthetic peptide derived from <i>Arabidopsis thaliana</i> HTA9 UniProt: <a href="#">Q9C944</a> , TAIR: <a href="#">At1g52740</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Immunogen affinity purified serum in PBS pH 7.4.
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µg
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Application information****Recommended dilution** 6 µg (ChIP), 1: 100 (IP), 1 : 1000 (WB)**Expected | apparent MW** 14.2 kDa (*Arabidopsis thaliana*)**Confirmed reactivity** *Arabidopsis thaliana***Not reactive in** *Nicotiana tabacum*, *Zea mays*

**Selected references**


[Bieluszewski](#) et al. (2022) NuA4 and H2A.Z control environmental responses and autotrophic growth in *Arabidopsis*. Nat Commun. 2022 Jan 12;13(1):277. doi: 10.1038/s41467-021-27882-5. PMID: 35022409; PMCID: PMC8755797.

[Bieluszewski](#) et al. (2022) NuA4 and H2A.Z control environmental responses and autotrophic growth in *Arabidopsis*. Nat Commun. 2022 Jan 12;13(1):277. doi: 10.1038/s41467-021-27882-5. PMID: 35022409; PMCID: PMC8755797.

[Kralemann](#) et al. (2020). Removal of H2Aub1 by ubiquitin-specific proteases 12 and 13 is required for stable Polycomb-mediated gene repression in *Arabidopsis*. Genome Biol. 2020 Jun 16;21(1):144. doi: 10.1186/s13059-020-02062-8.

[Gómez-Zambrano](#) et al. (2019). The repressive role of *Arabidopsis* H2A.Z in transcriptional regulation depends on AtBMI1 activity. Nat Commun. 2019 Jun 27;10(1):2828. doi: 10.1038/s41467-019-10773-1.

[Gómez-Zambrano](#) et al. (2018). *Arabidopsis* SWC4 Binds DNA and Recruits the SWR1 Complex to Modulate Histone H2A.Z Deposition at Key Regulatory Genes. Mol Plant. 2018 Mar 29. pii: S1674-2052(18)30122-9. doi: 10.1016/j.molp.2018.03.014.

**Application example**

1) *hta9, hta11* double mutant, about 1 µg protein; 2) mutant *hta9, hta11* about 2 µg protein; 3) wild-type about 2-3 µg protein; 4) wild-type about 4-5 µg protein from *Arabidopsis thaliana*.

Protein from *Arabidopsis thaliana* histone preparation (acid extracted, precipitated with acetone and resuspended in urea). Denatured with SDS-PAGE buffer at 90°C for 2 min, separated on 15% SDS-PAGE and blotted 1h to PVDF tank transfer. Blots were blocked with PBS + 3% BSA overnight at 4°C with agitation. Blot was incubated in the primary antibody at a dilution of 1:1 000 in PBS + 3% BSA for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed once with PBS-T for 5 min. Blot was incubated in secondary antibody PBS + 3% BSA (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera, AS09 602) diluted to 1:25 000 for 1h at RT with agitation. The blot was washed 3-4 times with PBS-T for 5 min. The blot was developed with Immobilon (Millipore) and exposed on an X-ray film for 1 min.