

Product no **AS09 514****Anti-HydA | Iron-hydrogenase HydA1/HydA2****Product information**

Immunogen	Recombinant, full length <i>Chlamydomonas reinhardtii</i> HydA-2 Q8VZZ0
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Additional information In *Chlamydomonas* **HydA** is present in low levels of **1 µg/liter of culture**. Therefore, an induction of cells by anaerobic adaptation or sulfur deprivation (10 x higher amount than with anaerobic adaptation) is necessary for successful detection using this antibody. Methods of HydA induction are described in [Hemschemeier et al. 2009](#).

To detect HydA1/A2 in *Chlamydomonas* extracts amount loaded per well corresponds to 2 µg of chlorophyll for sulfur deprived cells, where relatively much HydA1 is synthesized or corresponds to 2-4 µg of artificially anaerobic induced cultures, where the HydA1 protein level is lower. This antibody is recognizing 1 ng of recombinant HydA protein.

Application information

Recommended dilution	1 : 5000 (WB)
Expected apparent MW	53.7 48 kDa (after transit peptide is cleaved)
Confirmed reactivity	<i>Chlamydomonas reinhardtii</i>
Predicted reactivity	<i>Ostreococcus</i> sp.
	Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	HydA1 (497aa) has a calculated MW of 53.1 kDa, but this is including the signal peptide, which gets cleaved off. The protein without TP has a calculated MW of 47.5 kDa and runs according to its size at about 48 kDa.
	HydA2 (505aa) has a calculated MW of 53.7 kDa, but this is including the signal peptide, which gets cleaved off. The protein without TP can only be estimated, since the cleavage site is known only from in silico analysis. It has a calculated MW of 47.3 kDa and should run in the gel also according to its size.
	This antibody is binding recombinant HydA1/2 protein.

Selected references

[Nagy et al. \(2023\)](#). Photoautotrophic and sustained H₂ production by the pgr5 mutant of *Chlamydomonas reinhardtii* in simulated daily light conditions. *International Journal of Hydrogen Energy* Volume 53, 31 January 2024, Pages 760-769.

[Jokei et al. \(2020\)](#). Elimination of the flavodiiron electron sink facilitates long-term H₂ photoproduction in green algae. *Biotechnol Biofuels*. 2019 Dec 5;12:280. doi: 10.1186/s13068-019-1618-1.

[Lindblad et al. \(2019\)](#). CyanoFactory, a European consortium to develop technologies needed to advance cyanobacteria as chassis for production of chemicals and fuels. *Algal Research*, Volume 41, August 2019, 101510.

[Weiner et al \(2018\)](#), Overcoming the expression barrier of the ferredoxin hydrogenase chimera in *Chlamydomonas reinhardtii* supports a linear increment in photosynthetic hydrogen output. *Algal Research* Volume 33, July 2018, Pages 310-315

[Wei et al. \(2017\)](#). Light Intensity is Important for Hydrogen Production in NaHSO₃-Treated *Chlamydomonas reinhardtii*. *Plant Cell Physiol*. 2017 Mar 1;58(3):451-457. doi: 10.1093/pcp/pcw216.

[Eilenberg et al. \(2016\)](#). The dual effect of a ferredoxin-hydrogenase fusion protein in vivo: successful divergence of the photosynthetic electron flux towards hydrogen production and elevated oxygen tolerance. *Biotechnol Biofuels*. 2016 Aug 30;9(1):182. doi: 10.1186/s13068-016-0601-3. eCollection 2016.

[Liran et al. \(2016\)](#). Microoxic Niches within the Thylakoid Stroma of Air-Grown *Chlamydomonas reinhardtii* Protect [FeFe]-Hydrogenase and Support Hydrogen Production under Fully Aerobic Environment. *Plant Physiol*. 2016 Sep;172(1):264-71. doi: 10.1104/pp.16.01063. Epub 2016 Jul 21.


[Reifschneider-Wegner et al. \(2014\)](#). Expression of the [FeFe] hydrogenase in the chloroplast of *Chlamydomonas*

reinhardtii. *Int J of Hydrogen Energy*, Volume 39, Issue 8, 6 March 2014, Pages 3657–3665.

[Pinto et al. \(2013\)](#). Rubisco mutants of *Chlamydomonas reinhardtii* enhance photosynthetic hydrogen production. *Appl Microbiol Biotechnol*. May 7.

[Magneschi et al. \(2012\)](#). A Mutant in the ADH1 Gene of *Chlamydomonas reinhardtii* Elicits Metabolic 2 Restructuring during Anaerobiosis. *Plant Physiol*. January 23 (ahead of print).

Application example



50ng of purified protein (HydA1 and HydA2) were separated on 10% SDS-PAGE and blotted 25 min to PVDF membrane. Filters were blocked 1h with 3% low-fat milk powder in PBS-T (0.1% TWEEN 20) and probed with anti-HydA1/2 (AS09 514, 1:5000, over night at 4°C) and secondary anti-rabbit (1:10 000, 1 h) antibody (HRP conjugated, manufacture Pierce) in PBS-T containing 3% low fat milk powder. Antibody incubations were followed by washings in PBS-T (10, +10min and PBS (+5, +5 min). All washing steps were performed at RT with agitation. Signal was detected with ECL (Millipore) using CCD camera. Exposure time was 20 min.

The heterolog expressed proteins have both calculated MWs of 51 kDa (due to the tag) and run according to their size.

Courtesy Dr. Thomas Happe, Ruhr-Univer, Germany