

Product no **AS09 421****Anti-IAA | indole-3-acetic acid (N1) (for immunolocalization)****Product information**

<b>Immunogen</b>	BSA-conjugated, via N1 group of indole, indole-3-acetic acid (N1)
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Total IgG. Protein G purified in PBS.
<b>Format</b>	Liquid
<b>Quantity</b>	200 µg
<b>Storage</b>	Freeze upon arrival and store at -20°C; make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Additional information** | IAA | indole-3-acetic acid (N1) (for immunolocalization)**Application information**

<b>Recommended dilution</b>	Specific information about dilution is going to be included on the vial
<b>Confirmed reactivity</b>	indole-3-acetic acid (N1) tested so far in: <i>Eucalyptus globulus</i> , <i>Juncus prismatocarpus subsp. leschenaultii</i> Kirschner ( <i>J. prismatocarpus</i> ), <i>Juncus wallichianus</i> Laharpe ( <i>J. wallichianus</i> ), <i>Pinus radiata</i> , <i>Vitis vinifera</i>
<b>Predicted reactivity</b>	Indole-3-acetic acid (N1)
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Additional information</b>	Antibody can be used for direct detection of free IAA (see reference below). It does not bind IAA conjugates.  Antibody is provided in 50% glycerol. For larger quantity (1 mg)- please <a href="#">inquire</a> .  Steedman's wax embedding technique is recommended to be used with this antibody. The most critical issue is to keep the temperature below 37°C during the whole embedding procedure. More information can be found in <a href="#">Vitha et al. (2000)</a> .
<b>Selected references</b>	<a href="#">Nukazuka et al. (2021)</a> . A Role for Auxin in Triggering Lamina Outgrowth of Unifacial Leaves. <i>Plant Physiol.</i> 2021 Feb 23:kiab087. doi: 10.1093/plphys/kiab087. Epub ahead of print. PMID: 33620494. <a href="#">Dinis et al. (2018)</a> . Kaolin modulates ABA and IAA dynamics and physiology of grapevine under Mediterranean summer stress. <i>J Plant Physiol.</i> 2018 Jan;220:181-192. doi: 10.1016/j.jplph.2017.11.007. <a href="#">Escandón et al. (2016)</a> . Integrated physiological and hormonal profile of heat-induced thermotolerance in <i>Pinus radiata</i> . <i>Tree Physiol.</i> 2016 Jan;36(1):63-77. doi: 10.1093/treephys/tpv127. Epub 2016 Jan 12. <a href="#">Jesus et al. (2015)</a> . Salicylic acid application modulates physiological and hormonal changes in <i>Eucalyptus globulus</i> under water deficit. <i>Environ and Exp Botany</i> , Volume 118, October 2015, Pages 56-66. <a href="#">De Diego et al. (2013)</a> . Immunolocalization of IAA and ABA in roots and needles of <i>Pinus radiata</i> during drought and rewatering. <i>Tree Physiol.</i> , May; 33(5):537-549.