


Product no **AS09 513****Anti-Lhcb5 | CP26 (Lhcb5) homolog (Ostreococcus tauri)****Product information**

<b>Immunogen</b>	KLH-conjugated synthetic peptide derived from Lhcb4 (CP29) protein sequence from <i>Ostreococcus taurii</i> <a href="#">Q3B9U2</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	200 µl
<b>Reconstitution</b>	For reconstitution add 200 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Application information**

<b>Recommended dilution</b>	1 : 2000 (WB)
<b>Expected   apparent MW</b>	33 kDa
<b>Confirmed reactivity</b>	<i>Coccomyxa subellipsoidea</i> , <i>Ostreococcus tauri</i>
<b>Predicted reactivity</b>	Algae, <i>Ostreococcus lucimarinus</i>
	Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known

**Application example**

1.0-4.5 µg of chlorophyll from *Coccomyxa subellipsoidea* cells were loaded to lanes. Samples were denatured with Laemmli buffer at 75°C for 5 min and were separated on 12% SDS-PAGE and blotted 30 min to PVDF using wet transfer. Blot was blocked with 5% milk in TBS for 2h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1 : 2000 overnight at 4°C with agitation in 1% milk in TBS-T. The antibody solution was decanted and the blot was washed 4 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera, [AS09 602](#)) diluted to 1:25 000 in 1% milk in TBS-T for 1h at RT with agitation. The blot was washed 5 times for 5 min in TBS-T and 2 times for 5 min in TBS, and developed for 1 min with 1.25 mM luminol, 0.198 mM coumaric acid and 0.009% H<sub>2</sub>O<sub>2</sub> in 0.1 M Tris- HCl, pH 8.5. Exposure time in ChemiDoc System was 115 seconds.

Courtesy MSc Paweł Rogowski, Warsaw University, Poland