

Product no **AS14 2819****LHCSR1****Product information**

Immunogen	KLH-conjugated synthetic peptide derived from LHCSR1 protein sequence from <i>Chlamydomonas reinhardtii</i> , UniProt: P93664
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	27 kDa
Confirmed reactivity	<i>Bryopsis corticulans</i> , <i>Chlamydomonas reinhardtii</i> , <i>Nannochloropsis gaditana</i>
Predicted reactivity	Species of your interest not listed? Contact us
Not reactive in	<i>Lobosphaera incisa</i> , <i>Phaeodactylum tricornutum</i>
Additional information	This antibody is also recognizing recombinant LHCSR1 overexpressed in <i>E.coli</i> as described in Perozeni et al. (2020) .
Selected references	<p>McQuillan et al. (2023). Proteomic characterization of a lutein-hyperaccumulating <i>Chlamydomonas reinhardtii</i> mutant reveals photoprotection-related factors as targets for increasing cellular carotenoid content. <i>Biotechnol Biofuels Bioprod</i> . 2023 Nov 4;16(1):166. doi: 10.1186/s13068-023-02421-0.</p> <p>Cazzaniga et al. (2022). Engineering astaxanthin accumulation reduces photoinhibition and increases biomass productivity under high light in <i>Chlamydomonas reinhardtii</i>. <i>Biotechnol Biofuels Bioprod</i>. 2022 Jul 11;15(1):77. doi: 10.1186/s13068-022-02173-3. PMID: 35820961; PMCID: PMC9277849.</p> <p>Redekop et al. (2020). PsbS Contributes to Photoprotection in <i>Chlamydomonas Reinhardtii</i> Independently of Energy Dissipation . <i>Biochim Biophys Acta Bioenerg</i> . 2020 Jun 1;1861(5-6):148183.doi: 10.1016</p> <p>Lammermann et al. (2020). Ubiquitin ligase component LRS1 and transcription factor CrHy5 act as a light switch for photoprotection in <i>Chlamydomonas</i>. doi.org/10.1101/2020.02.10.942334 bioRxiv</p> <p>Roach et al. (2020). The non-photochemical quenching protein LHCSR3 prevents oxygen-dependent photoinhibition in <i>Chlamydomonas reinhardtii</i>. <i>J Exp Bot</i>. 2020 Jan 16. pii: eraa022. doi: 10.1093/jxb/eraa022.</p> <p>Gabilly et al. (2019). Regulation of photoprotection gene expression in <i>Chlamydomonas</i> by a putative E3 ubiquitin ligase complex and a homolog of CONSTANS. <i>Proc Natl Acad Sci U S A</i>. 2019 Aug 12. pii: 201821689. doi: 10.1073/pnas.1821689116.</p>

Application example

10 µg of a total cell extract of *Chlamydomonas reinhardtii*: WT_P_HL_High CO₂ (1), WT_P_HL (2), Npq4_P_HL (3), were separated on Bolt® 4-12% Bis-Tris Plus Gels (precast) and blotted to PVDF using Bolt® transfer system for 1h. Blots were blocked with 5% BSA/milk for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1 000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and

3 times for 5 min in PBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, [AS09 602](#), Agrisera) diluted to 1:10 000 in for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL according to the manufacturer's instructions. Exposure time was 10 seconds.

P: photoautotrophically grown cells

HL: High light illumination(light intensity:500 μ E)

Courtesy Dr. Roberta Croce, Biophysics of Photosynthesis Dep. Physics and Astronomy Faculty of Sciences VU University Amsterdam, The Netherlands