

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS12 2104

Anti-LSD1 | Lesion simulating disease 1 (chicken antibody)

Product information

Immunogen KLH-conjugated synthetic peptide derived from *Arabidopsis thaliana* LSD1 sequence, UniProt: <u>P94077</u>, TAIR:

AT4G20380

Host Chicken

Clonality Polyclonal

Purity Immunogen affinity purified IgY in PBS pH 7.4. Contains 0.02% sodium azide.

Format Liquid

Quantity 50 μg

Storage Store at 4°C; make aliquots to avoid working with a stock. Please remember to spin the tubes briefly prior to opening

them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution 1:1000 (WB)

Expected | apparent

MW 19.5 KI


Predicted reactivity Brassica olereacea, Pisum sativum

Species of your interest not listed? Contact us

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Additional information This antibody has so far confirmed reactivity on recombinant LSD1

Application example



50 g of total protein from *Arabidopsis thaliana* wilde type, Isd1 mutant and transgenic plants overexpressin LSD1-GFP protein under 35S promotor were extracted with following extraction buffer: 0.0625 M Tris-HCl pH 6.8, 1 % SDS, 10 % glycerol, 0.01% beta-mercaptoethanol) were separated in 10 % SDS-PAGE and blotted for 16 hours to PVDF at 4 °C using tank transfer. Blots were blocked in PBS-T with 4.5 % non-fat milk for 4 hours at room temperature with agitation. Blot was incubated in the primary antibody at a dilution of 1.500 for 20 hours at 4 °C with agitation. The antibody solution was decanted and the blot was washed 5x for 10 minutes in PBS-T at RT with agitation. Blot was incubated in secondary antibody, anti-chicken IgY, HRP conjugated diluted to 1:7500 in PBS for 1 hour at RT with agitation. The blot was washed as above and developed for 5 minutes with ECL according to the manufacture's instruction. Exposure time was 1 minute.

Courtesy of Dr. Keun Pyo Lee, SIBS, China