


Product no **AS12 2633****MPK6 | Mitogen-activated protein kinase 6****Product information**

| | |
|-----------------------|---|
| Immunogen | KLH-conjugated synthetic peptide derived from <i>Arabidopsis thaliana</i> MPK6 protein, UniProt: Q39026 , TAIR: At2g43790 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Purity | Immunogen affinity purified serum in PBS pH 7.4. |
| Format | Lyophilized |
| Quantity | 50 µg |
| Reconstitution | For reconstitution add 50 µl of sterile water |
| Storage | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube. |

Application information

| | |
|-------------------------------|--|
| Recommended dilution | 1 : 1000 (WB) |
| Expected apparent MW | 45 kDa |
| Confirmed reactivity | <i>Arabidopsis thaliana</i> |
| Predicted reactivity | <i>Brassica napus</i> , <i>Gossypium mexicanum</i> Species of your interest not listed? Contact us |
| Not reactive in | <i>Lupinus luteus</i> , <i>Picea abies</i> , <i>Solanum tuberosum</i> |
| Selected references | Butler et al. (2019). Soybean resistance locus Rhg1 confers resistance to multiple cyst nematodes in diverse plant species. <i>Phytopathology</i> . 2019 Aug 12. doi: 10.1094/PHYTO-07-19-0225-R. Wang and Auwerx (2017). Systems Phytohormone Responses to Mitochondrial Proteotoxic Stress. <i>Mol Cell</i> . 2017 Nov 2;68(3):540-551.e5. doi: 10.1016/j.molcel.2017.10.006. |

Application example

0.2g of 21 days old *Arabidopsis thaliana* leaf tissue C=Col, 6= mpk6 (T-DNA mutant), 3= mpk3 (T-DNA mutant) was homogenized with 250 µl Buffer A (50 mM Tris – HCl pH 7.5, 0.33 M sucrose, 5 mM EDTA, 150 mM NaCl, complete protease inhibitor cocktail). The crude extracts were subjected to centrifugation at 10 000 g for 10 min to pellet the insoluble material. Material was extracted with 250 µl of Buffer A (50 mM Tris – HCl pH 7.5, 0.33 M sucrose, 5 mM EDTA, 150 mM NaCl, complete protease inhibitor cocktail). The crude extracts were subjected to centrifugation at 10 000 g for 10 min to pellet the insoluble material. Supernatant (120 µL) + 4XSDS sample buffer 40 µL = 200 µL samples 95 °C 5min 10 µL were separated on 10 % SDS-PAGE and blotted 1h to PVDF using semi-dry transfer. Blots were blocked with skim milk for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody in TBS-T at a dilution of 1: 10 000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera [AS09 602](#)) diluted to 1:75 000 in for 1h at RT with agitation. The blot was washed as above and developed with chemiluminescence according to the manufacturer's instructions.

Courtesy of Chika Tateda, University of Chicago, USA