


Product no **AS11 1776****Anti-N-YFP | N-terminal of YFP****Product information**

Immunogen	KLH-conjugated synthetic peptide derived from N-terminal of YFP protein.
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 2000 (WB)
Confirmed reactivity	N-YFP tagged proteins from <i>Arabidopsis thaliana</i> , <i>Escherichia coli</i> , <i>Nicotiana tabacum</i>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	YFP molecular weight is 27 kDa, however detected band is going to have MW increased by a fused protein
Selected references	<p>Li et al. (2021) Two ubiquitin-associated ER proteins interact with COPT copper transporters and modulate their accumulation, <i>Plant Physiology</i>, 2021, kiab381, https://doi.org/10.1093/plphys/kiab381</p> <p>Lung et al. (2021) Oxylin signaling in salt-stressed soybean is modulated by ligand-dependent interaction of Class II acyl-CoA-binding proteins with lipoxygenase. <i>Plant Cell</i>. 2021 Dec 17:koab306. doi: 10.1093/plcell/koab306. Epub ahead of print. PMID: 34919703.</p> <p>Schultz-Larsen et al. (2018). The AMSH3 ESCRT-III-Associated Deubiquitinase Is Essential for Plant Immunity. <i>Cell Rep</i>. 2018 Nov 27;25(9):2329-2338.e5. doi: 10.1016/j.celrep.2018.11.011.</p>

Application example

Arabidopsis thaliana leaf lysate which is co-expressing test protein 1 with nYFP tags (~60 kDa) and test protein 2 with cYFP-tag (~40 kDa). In each case only one band is visible demonstrating that the antibodies do not cross-react. Sample preparation and immunoblot analysis were carried out as described in [Karnik et al.](#), *Plant Cell* 2015, March 2015 vol. 27 no. 3 675-694. For immunoblot analysis of plant tissues, leaves were excised and flash frozen in liquid N₂. Frozen tissue was ground in equal volumes (w/v) of homogenization buffer containing 500 mM sucrose, 10% glycerol, 20 mM EDTA, 20 mM MEGTA, Protease Inhibitor (Roche), 10 mM Ascorbic acid, 5 mM DTT, and 50 mM Tris-HCl, pH 7.4, and centrifuged at 13,000g and 4°C for 30 min to pellet debris. Supernatant was diluted 1:1 in 23 Laemmli buffer containing 2.5% 2-mercaptoethanol, heated to 95°C for 10 min, and separated by SDS-PAGE on a 12% Acrylamide gel. Ponceau S-stained Rubisco bands were used as loading standards for plant samples. Membrane type: Cellulose Nitrate (GE Healthcare) Blocking reagent: GE Healthcare Wash buffer: Tris Buffered Saline, 0.5% Tween Exposure time: 10 – 20 seconds.

Courtesy of Dr. Rucha Karnik, University of Glasgow, UK