


Product no **AS12 2612****Anti-NRT2,1 | Nitrate transporter 2,1****Product information**

<b>Immunogen</b>	KLH-conjugated synthetic peptide derived from <i>Arabidopsis thaliana</i> NRT2.1 C-terminal sequence, UniProt: <a href="#">O82811</a> , TAIR: <a href="#">AT1G08090</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Immunogen affinity purified serum in PBS pH 7.4.
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µg
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Application information**

<b>Recommended dilution</b>	1 : 5000 (WB)
<b>Expected   apparent MW</b>	57.7   45 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i>
<b>Predicted reactivity</b>	<i>Brassica juncea</i> , <i>Brassica napus</i> , <i>Brassica rapa</i> , <i>Ricinus communis</i> Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	<i>Alexandrium pacificum</i> , <i>Glycine max</i> , <i>Solanum tuberosum</i>
<b>Additional information</b>	Microsomal fraction is recommended to be used as it will have higher amount of NRT2,1 protein, If a total protein extract is used SDS should be applied from the beginning in extraction buffer as well as protease inhibitor cocktail
<b>Selected references</b>	<a href="#">Zou et al. (2019)</a> . Phosphorylation at Ser28 stabilizes the Arabidopsis nitrate transporter NRT2.1 in response to nitrate limitation. <i>J Integr Plant Biol.</i> 2019 Jul 24. doi: 10.1111/jipb.12858.

**application information**

50 µg of microsomal protein from wild type *Arabidopsis thaliana* (Col 0) roots at various stages of development (21 and 42 days) and from NRT2.1 deletion mutant (KO-NRT2.1) were solubilized with Laemmli x2 sample buffer and were separated on 12 % SDS-PAGE gel. The proteins were transferred (tank transfer, 100 V, 1h) to a PVDF membrane (Immobilon-P, 0.45 µm, Millipore). Blot was blocked for 2h with agitation at room temperature in blocking buffer (SuperBlock, Thermo) containing 0.05 % Tween 20 (Thermo). Blot was then incubated for 2h at RT and with agitation in the fresh blocking buffer containing the primary antibody at a dilution of 1:5 000. The antibody solution was decanted and the blot was washed 3 times for 10 min in PBS-T (0.05 % Tween 20, Thermo) at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera, [AS09 602](#)) diluted to 1:10 000 in blocking buffer, for 2h at RT with agitation. The blot was washed 1 time for 10 min in PBS-T and 2 times for 10 min in PBS. The blot was then incubated with ECL reagent (Super Signal West Pico, Thermo) for 5 min and the chemiluminescence was recorded on LAS 3000, Fuji. Exposure times were between 5-15 min using standard (lowest) sensitivity.

Courtesy of Dr. Wojciech Szponarski, French National Institute of Agricultural Research, France