


Product no **AS15 2833****Anti-PAB | protein in chloroplast atpase biogenesis****Product information**

<b>Immunogen</b>	Recombinant PAB from <i>Zea mays</i> , <a href="#">GRMZM2G110258</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µl
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile water
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Application information**

<b>Recommended dilution</b>	1 : 1000 (WB)
<b>Expected   apparent MW</b>	35 kDa (without transit peptide)
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Zea mays</i>
<b>Predicted reactivity</b>	<i>Brachypodium distachyon</i> , <i>Gossypium raimondii</i> , <i>Hordeum vulgare</i> , <i>Oryza sp.</i> , <i>Saccharum hybrid cultivar R570</i> , <i>Setaria italica</i> , <i>Sorghum bicolor</i> , <i>Theobroma cacao</i> , <i>Triticum aestivum</i> Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known

**application information**

Lanes contain ~5 µg of total protein from *Zea mays* seedling leaves (upper image), *Arabidopsis thaliana* seedling leaves (lower image), or isolated chloroplast stroma. Leaf extract from homozygous *pab* mutants (hypomorphic alleles), and purified recombinant antigen were analyzed as controls. Proteins were separated on 12% SDS-PAGE and blotted overnight to nitrocellulose using tank transfer. Blots were blocked for 45 min at room temperature (RT) in TBST +4% dry milk with agitation. The blot was incubated in the primary antibody at a dilution of 1: 1 000 for 2h at RT with agitation. The antibody solution was decanted and the blot was rinsed twice in TBST, then washed three times for 10 min in TBST at RT with agitation. The blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera, [AS09 602](#)) diluted to 1:10 000 in TBST +1% dry milk for 1h at RT with agitation. The blot was washed as above and incubated for ~2 minutes in ECL reagents prior to imaging with a LiCor digital imager. Exposure time was ~30 seconds.



This product is **for research use only** (not for diagnostic or therapeutic use)

**contact: [support@agrisera.com](mailto:support@agrisera.com)**

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

Courtesy of Dr. Alison Barkan, The University of Oregon, USA