

Product no **AS06 116****Anti-PGL35 | Plastoglobulin 35; FIB1a; FBN1a****Product information**

Immunogen	Recombinant <i>Arabidopsis thaliana</i> PGL35 protein O81439 , At4g04020
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	200 µl
Reconstitution	For reconstitution add 200 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
Additional information	Cellular [compartment marker] of chloroplast plastoglobules. For IC samples were embedded in Lowicryl HM20 and sectioned into 100-nm-thick sections and placed on Formvar-coated gold slot grids. The sections were blocked for 20 min with a 5% (w/v) solution of nonfat milk in TBS plus 0.1% Tween 20 (TBST). Anti-PGL antibodies were diluted 1:20 in a solution of 2.5% nonfat milk in TBST at room temperature for 1 h. The sections were rinsed in a stream of TBS plus 0.5% Tween 20 and then transferred to the secondary antibody (anti-rabbit IgG 1:20 in TBST) conjugated to 10-nm gold particles for 1 h. Images of localization can be found in Austin et al. (2006).

Application information

Recommended dilution	1 : 20 (IC), 1 : 1000-1 : 3000 (WB)
Expected apparent MW	35 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Citrus reticulata</i> , <i>Gossypium hirsutum</i> cv. Deltapine 90
Predicted reactivity	<i>Brassica napus</i> , <i>Brassica campestris</i> , <i>Capsicum annum</i> , <i>Hordeum vulgare</i>
	Species of your interest not listed? Contact us
Not reactive in	<i>Chlamydomonas reinhardtii</i> , <i>Pheodactylum tricornutum</i> , <i>Pisum sativum</i>
Additional information	AtPGL35 is highly similar to <i>Pisum sativum</i> PG1
Selected references	(2021) . Autophagy is required for lipid homeostasis during dark-induced senescence. <i>Plant Physiology</i> , 2021;, k1aa120 Luo et al. (2015) . Distinct carotenoid and flavonoid accumulation in a spontaneous mutant of Ponkan (<i>Citrus reticulata</i> Blanco) results in yellowish fruit and enhanced postharvest resistance. <i>J Agric Food Chem</i> . 2015 Sep 2. Gámez-Arjona et al. (2014) . Starch synthase 4 is located in the thylakoid membrane and interacts with plastoglobule-associated proteins in <i>Arabidopsis</i> . <i>Plant J</i> . 2014 Oct;80(2):305-16. doi: 10.1111/tj.12633.