

This product is for research use only (not for diagnostic or therapeutic use)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS14 2785 Anti-PRP40B | pre-mRNA-processing protein 40B

Product information

 Immunogen
 Recombinant Arabidopsis thaliana PRP40b, UniProt:F4JCC1, TAIR: At3g19670

 Host
 Rabbit

 Clonality
 Polyclonal

 Purity
 Immunogen affinity purified serum in PBS pH 7.4.

 Format
 Lyophilized

 Quantity
 50 µg

 Reconstitution
 For reconstitution add 50 µl of sterile water

 Storage
 Store Iyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 2000 (WB)
Expected apparent MW	113 kDa
Confirmed reactivity	Arabidopsis thaliana
Predicted reactivity	Brassica rapa
	Species of your interest not listed? Contact us
Not reactive in	

Application example



Total protein from rosette leaves, 30 ug of *Arabidopsis thaliana* whole leaf extract (prp40ab) **(1)**, 30 ug of *Arabidopsis thaliana* whole leaf extract (wild-type) **(2)**, extracted with buffer (100mM Tris, 10% glicerol, 5mM EDTA, 5mM EGTA, 0.15M NaCl, 0.75% Triton X100, 0.05% SDS, 1mM DTT) and denatured with 0.06M Tris-HCl, 5% glycerol, 2% SDS, 4% -mercaptoethanol, 0.0025% bromophenol blue at 95 C for 5 min were separated on 8 % SDS-PAGE and blotted 1h to PVDF using semi-dry. Blots were blocked with 5% low-fat dried milk in TBS + 0,1%Tween for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 2 000 for 1h at RT with agitation in 2% low-fat dried milk in TBS + 0,1%Tween. The antibody solution was decanted and the blot was rinsed briefly twice, then washed 3 times for 10 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from) diluted to 1:10 000 in 2% low-fat dried milk in TBS + 0,1%Tween for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL of high sensitivity. Exposure time was 60 seconds.

MW markers: PageRuler Plus Prestained (ThermoFisher)

Courtesy of Agata Stępień, Adam Mickiewicz University, Poland

1/1