

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS10 1524

Rabbit anti-Rat IgG (H&L), HRP conjugated

Product information

Immunogen Purified rat IgG (H&L)

Host Rabbit

Clonality Polyclonal

Purity Immunogen affinity purified rabbit IgG.

Format Lyophilized

Quantity 1 mg

Reconstitution

For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily

Storage

Store lyophilized material at 2-8 °C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20 °C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20 °C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.

Additional information

Purity of this preparation is > 95% based on SDS-PAGE. Antibody concentration is 1.0 mg/ml. Antibody is supplied in 10 mM sodium phosphate, 0.15 M sodium chloride, pH 7.2.1 % (w/v) B, Protease/IgG free. Contains 0.1 % (v/v) Kathon CG as preservative of bacterial growth.

Based on immunoelectrophoresis, this antibody reacts with: heavy chains on rat lgG, light chains on all rat immunoglobulins. Based on immunoelectrophoresis, no reactivity is observed to: non-immunoglobulin rat serum proteins

Application information

Recommended dilution

This conjugate is suitable for all immunoassay applications, The optimal working dilution should be determined by the investigator

Selected references

Quian et al. (2015). Bone Marrow-Derived Mesenchymal Stem Cells Repair Necrotic Pancreatic Tissue and Promote Angiogenesis by Secreting Cellular Growth Factors Involved in the SDF-1 /CXCR4 Axis in Rats. Stem Cells International Volume 2015 (2015), Article ID 306836.