


Product no **AS16 3148****Anti-SAM1-4 | S-adenosylmethionine synthase****Product information**

Immunogen	KLH-conjugated peptide derived from protein sequence of <i>Arabidopsis thaliana</i> SAM1-4, UniProt: P23686 , P17562 , Q9SJL8 , Q9LUT2 , TAIR: At1g02500 , At4g01850 , At3g17390 , At2g36880
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.


Application information

Recommended dilution	1 : 3000 (WB)
Expected apparent MW	43.2 45 kDa (<i>Arabidopsis thaliana</i>)
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Malus domestica</i>
Predicted reactivity	<i>Beta vulgaris</i> , <i>Brassica sp.</i> , <i>Citrus sp.</i> , <i>Coffea canephora</i> , <i>Caomelina sativa</i> , <i>Capsella rubella</i> , <i>Cucumis melo</i> , <i>Cucumis sativus</i> , <i>Genlisea aurea</i> , <i>Gentiana triflora</i> , <i>Guzmania wittmacki</i> , <i>Gossypium raimondii</i> , <i>Eucalyptus grandis</i> , <i>Eutrema salsugineum</i> , <i>Ipomoea batatas</i> , <i>Jatropha curcas</i> , <i>Musa acuminata</i> , <i>Nicotiana sp.</i> , <i>Oryza brachyantha</i> , <i>Phoenix dactylifera</i> , <i>Populus sp.</i> , <i>Prunus sp.</i> , <i>Ricinus communis</i> , <i>Sesamum indicum</i> , <i>Setaria italica</i> , <i>Solanum pennellii</i> , <i>Solanum tuberosum</i> , <i>Spinacia oleracea</i> , <i>Tarenaya hassleriana</i> , <i>Theobroma cacao</i> , <i>Zea mays</i>
	Species of your interest not listed? Contact us
Not reactive in	<i>Prunus domestica</i>

Application example

70 µg of total protein from *Arabidopsis thaliana* wt Col-0, *SAM1-4* knockdown, and two *oeSAM3-YFP* lines extracted with mortar and pestle using 2xSDS loading buffer (100 mM Tris-HCl pH 6.8, 4% SDS, 0.02% bromophenol blue, 200 mM DTT), and denatured in the same buffer at 95°C for 10 min. Samples were separated on 11% SDS-PAGE and blotted for 1h to PVDF using tank transfer. Blots were blocked with 5% milk powder in TBS-T overnight in a cold room with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1 000 for 1h at RT with agitation. The antibody solution was decanted and the blot was rinsed five times for 4 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG HRP-conjugated, from Agrisera) diluted to 1: 20 000 in for 2h at RT with agitation. The blot was washed as above and developed for 2 min with chemiluminescent detection reagent. The proteins were detected using CCD Image Fusion Fx7 after 60 seconds exposure time.

Courtesy of Louis-Valentin Meteignier, LGBP, Faculté des Sciences de Luminy, France



Leaf tissue of *Arabidopsis thaliana* Col-8 plants was disrupted via mortar and pestle under liquid nitrogen. Leaf protein was extracted by addition of appropriate amounts of extraction buffer (6 M Guanidine-HCl, 100 mM HEPES, 5 mM EDTA and 1x Halt™ Protease Inhibitor-Cocktail). Extracted protein was chloroform/methanol-precipitated and resolved in 2 % SDS, 50 mM Tris (pH 7.4). Protein concentration was determined using BCA assay. 100 µg of cleaned protein were mixed with 4x LDS sample buffer (final conc. 1x), 50 mM DTT (final) and heated 2x to 95 °C for 5 minutes for denaturation. 2.5 µg of protein were loaded per lane on a 12 % SDS PAGE gel. After separation, proteins were blotted to a PVDF membrane for 30 minutes with 25 V and 1 A in a semi-dry blot. Blots were blocked with 5 % Milk in TBS (= Tris buffered saline) for 2 h at room temperature. Blots were subsequently washed with TBS-T (TBS + 0.05 % Tween20) for 1h at room temperature with 3 exchanges of wash buffer. Primary antibody (anti SAM1-4) was used at a dilution of 1:2000 in 1 % Milk in TBS-T and blots were incubated over night at 4 °C with gentle agitation. Primary antibody was decanted and blots were again washed for 1 h with TBS-T as mentioned above. Secondary antibody (anti Rabbit IgG, HRP conjugated) was used at a dilution of 1:20000 in 1 % Milk in TBS-T and blots were incubated for 1 h at room temperature. After decanting of secondary antibody, blots were again washed with TBS-T for 1h. Before Chemiluminescence detection, blots were shortly washed with TBS without Tween20 3x 5 minutes at room temperature. Chemiluminescent detection reagent was used for signal detection. Images of the blots were obtained using ChemiDoc™ XRS (Bio-rad) in high resolution mode. Exposure time was 20 seconds.

Courtesy of phd student Andreas Perrar, Prof. Dr. Pitter Huesgen group, Forschungszentrum Jülich, Germany