

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS10 1333**
Cystatin C

Product information

Background	Cystatin C (or cystatin 3, called also gamma trace, post-gamma-globulin or neuroendocrine basic peptide) is a protein which can be found in normal human urine and is used as a biomarker of kidney function.
Immunogen	human cystatin C CAA36497
Host	Goat
Clonality	Polyclonal
Purity	Affinity purified
Format	Liquid
Quantity	1 mg
Storage	Store at 2-8°C.
Tested applications	Immunoassays

Application information

Recommended dilution	The optimal working dilution should be determined by the investigator.
Expected apparent MW	16 14 kDa
Confirmed reactivity	Human cystatin C
Predicted reactivity	Cat, Mouse, Rat
Not reactive in	No confirmed exceptions from predicted reactivity are currently known.
Additional information	Affinity purified using solid phase human Cystatin C. Antibody concentration is > 4.5 mg/ml (E 1% at 280 nm = 13.0). Antibody is supplied in 10 mM sodium phosphate, 0.15 M sodium chloride, pH 7.2. Contains 0.05% (w/v) sodium azide as preservative. Antibody purity is > 95% based on SDS-PAGE. For high resolution images, please visit the specific product page at www.agrisera.com

For high resolution images, please visit the specific product page at www.agrisera.com