

# Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

Product no **AS16 3981**

## 2b protein [Cucumber mosaic virus]


### Product information

<b>Immunogen</b>	Recombinant 2b protein [Cucumber mosaic virus] Protein accession number: <a href="#">NP_619631</a> .
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µl
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile water.
<b>Storage</b>	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
<b>Additional information</b>	This product can be sold containing proclin if requested

### Application information

<b>Recommended dilution</b>	1 : 5000 (WB)
<b>Expected   apparent MW</b>	12.7   18 kDa
<b>Confirmed reactivity</b>	2b protein [Cucumber mosaic virus]
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known.
<b>Selected references</b>	<a href="#">Wu et al. (2020)</a> . WUSCHEL triggers innate antiviral immunity in plant stem cells. <i>Science</i> . 2020 Oct 9;370(6513):227-231. doi: 10.1126/science.abb7360. PMID: 33033220. <a href="#">Nemes et al. (2019)</a> . Symptom recovery is affected by Cucumber mosaic virus coat protein phosphorylation. <i>Virology</i> Volume 536, October 2019, Pages 68-77. <a href="#">Zhang et al (2006)</a> . Cucumber mosaic virus-encoded 2b suppressor inhibits Arabidopsis Argonaute1 cleavage activity to counter plant defense. <i>Genes &amp; development</i> 20: 3255-3268.

#### Application example



80 µg of *Arabidopsis thaliana* soluble total cell extract from FNY 2b transgenic plants or the indicated amounts of the untagged recombinant proteins (extracted in 2 x SDS buffer (0.125M Tris pH 6.8, 4% (w/v) SDS, 20%(v/v) glycerol, 0.2M DTT, 0.02% bromophenol blue)) was separated on 15% SDS-PAGE and blotted 1h to PVDF membrane. Filters were blocked 1h with 5% low-fat milk powder in PBS-T (1 X PBS buffer; 0.5% TWEEN20) and probed with the serum of anti-FNY 2b antibody (1:5 000, 1h) and secondary anti-rabbit (1:5000, 1 h) antibody (HRP conjugated) in PBS-T containing 5% low fat milk powder. Antibody incubations were followed by washings in PBS-T. All steps were performed at RT with agitation. Blots were developed for 5 min with ECL-Prime detection reagent according the manufacturer's instructions. Exposure time was 20 seconds.

# Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

Courtesy of Dr. Xiuren Zhang, Texas A&M University, USA