

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS09 591**

AtpH | ATP synthase subunit c (chloroplastic)

Product information

Immunogen	KLH-conjugated peptides derived from AtpH subunit c of <i>Arabidopsis thaliana</i> UniProt: P56760 , TAIR: AtCg00140 and <i>Chlamydomonas reinhardtii</i> UniProt: Q37304
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	100 µl
Reconstitution	For reconstitution add 100 µl of sterile water.
Storage	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
Additional information	This product can be sold containing ProClin if requested.

Application information

Recommended dilution	1 : 10 000 (WB)
Expected apparent MW	8 kDa (for <i>Arabidopsis thaliana</i>)
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Chlamydomonas reinhardtii</i>
Predicted reactivity	Algae, <i>Cannabis sativa</i> , <i>Glycine max</i> , <i>Hordeum vulgare</i> , <i>Oryza sativa</i> , <i>Ostreococcus tauri</i> , <i>Physcomitrella patens</i> , <i>Pinus thunbergii</i> , <i>Pisum sativum</i> , <i>Populus alba</i> , <i>Zea mays</i> , <i>Vitis vinifera</i> Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known.
Additional information	Please note that increased incubation at 95°C (20-30 min) prior to loading is recommended to break the multimeric c-mer structure, detection of partial ring structures (e.g. 5 or 6 subunits) may occur. For high resolution images, please visit the specific product page at www.agrisera.com
Selected references	Schulz et al. (2017) . Molecular architecture of the N-type ATPase rotor ring from <i>Burkholderia pseudomallei</i> . EMBO Rep. 2017 Apr;18(4):526-535. doi: 10.15252/embr.201643374.

Application example

Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com



10 ug of chlorophyll/well of *Chlamydomonas reinhardtii* total cell extract (**1**), *Chlamydomonas reinhardtii* subunit gamma deletion mutant thylakoid membrane fraction (**2**), *Arabidopsis thaliana* thylakoid membrane fraction (**3**), *Chlamydomonas reinhardtii* thylakoid membrane preparation (4) were separated on 12-18% acrylamide-8M urea gel and blotted to nitrocellulose membrane. Filters were blocked 1 h with 5% dry milk in 1 x PBS and probed with anti-ATP synthase subunit c antibody (**AS09 591**, 1: 10 000, 1h) and secondary HRP-conjugated anti-rabbit antibody (1: 10 000, 1 h) in 1 x PBS containing 5% dry milk. All steps were performed at RT with agitation. Signal was detected with chemiluminescent detection, exposure time 30".

Arabidopsis membrane preparation has been done according to [Lezhneva et al. \(2008\)](#) A novel pathway of cytochrome c biogenesis is involved in the assembly of the cytochrome b6f complex in arabidopsis chloroplasts. J Biol. Chem., 283:24608-24616 and *Chlamydomonas* membranes were prepared according to [Chua & Bennoun \(1975\)](#) Thylakoid membrane polypeptides of *Chlamydomonas reinhardtii*: wild-type and mutant strains deficient in photosystem II reaction center. PNAS 72:2175-2179

Courtesy Dr. Yves Choquet, CNRS, France