

Product no **AS16 3353**

**Goat anti-Human IgG Fc, F(ab)'2 Fragment, DyLight® 488 conjugated, min, cross-reactivity to human IgA, IgM or IgG F(ab)'2 fragment**

## Product information

**Immunogen** | Purified Human IgG, Fc fragment

**Host** | Goat

**Clonality** | Polyclonal

**Purity** | Immunogen affinity purified goat IgG.

**Format** | Lyophilized

**Quantity** | 0.5 mg

**Reconstitution** | For reconstitution add 0.55 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily

**Storage** | Store freeze-dried powder at 2-8 °C.

**Additional information** | Antibody purity is > 90% based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Human IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1% (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative.

Based on IEP, this antibody reacts with: • heavy ( ) chains on human IgG.  
Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgA or IgM.

## Application information

**Recommended dilution** | 1 : 20-1 : 2000 (Flow cyt), (IL)

**Confirmed reactivity** | Goat anti-Human IgG Fc, F(ab)'2 Fragment