

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS16 3689**
GUS | Beta-glucuronidase


Product information

| | |
|-------------------------------|--|
| Immunogen | KLH-conjugated synthetic peptide derived from <i>Escherichia coli</i> GUS protein at amino acid 358. |
| Host | Rabbit |
| Clonality | Polyclonal |
| Purity | Serum |
| Format | Lyophilized |
| Quantity | 50 µl |
| Reconstitution | For reconstitution add 50 µl of sterile water. |
| Storage | Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes. |
| Additional information | This antibody will not work with pCAMBIA vectors due to mutation starting at amino acid position 358. |

Application information

| | |
|-------------------------------|---|
| Recommended dilution | 1 : 10 000 (WB) |
| Expected apparent MW | depends upon a MW of a protein which is expressed with GUS |
| Confirmed reactivity | GUS |
| Not reactive in | No confirmed exceptions from predicted reactivity are currently known. |
| Selected references | Nizkorodova et al. (2020) . The Effect of Translation Promoting Site (TPS) on Protein Expression in E. coli Cells. Mol Biotechnol (2020). doi.org/10.1007/s12033-020-00251-1 Nizkorodova and Isakov (2019) . New insights into the mechanism of action of e-element enhancer in E.coli. Eurasian Journal of Applied Biotechnology 2/2019 |

Application example



Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

100 µg of total protein from *Arabidopsis thaliana* wild-type **(1)** and transgenic plants expressing GUS under HRT promoter **(2)** were extracted with RNeasy lysis buffer containing 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 10 mM MgCl₂, 5 mM EDTA, 10% glycerol, 0.1% NP-40, 0.1% protease inhibitor cocktail. The proteins were denatured by boiling for 5 min, separated on 10% SDS-PAGE and blotted 30 min to PVDF using tank transfer. Blots were blocked with TBST containing 5% non-fat dry milk for 1 h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1:10 000 for 2 h at RT with agitation in TBS-T. The antibody solution was decanted and the blot was rinsed briefly twice, then washed 3 times for 15 min each in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1:10 000 in TBS-T for 1 h at RT with agitation. The blot was washed as above and developed with Clarity MAX (Bio-RAD). Exposure time was 10 seconds.

Courtesy of Dr. Pradeep Kachroo, University of Kentucky, USA