

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS06 172**

PsaA | PSI-A core protein of photosystem I

Product information

Background	PsaA is a core protein of photosystem I. In plants and cyanobacteria, the primary step in oxygenic photosynthesis, the light induced charge separation, is driven by two large membrane intrinsic protein complexes, the photosystems I and II. Synonym: Photosystem I P700 chlorophyll a apoprotein A1.
Immunogen	N-terminal part of recombinant PsaA protein from <i>Chlamydomonas reinhardtii</i> P12154
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water.
Storage	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
Tested applications	Immunogold (IG), Western blot (WB), Blue Native PAGE (BN-PAGE)
Related products	Collection of antibodies to PSI proteins recommended secondary antibody Plant and algal protein extraction buffer Secondary antibodies
Additional information	PsaA is a hydrophobic protein and we recommend to use PVDF membrane for transfer to assure best results. This product can be sold containing ProClin if requested.

Application information

Recommended dilution	1 : 20 (IG), 1 : 1000-1 : 5000 (WB)
Expected apparent MW	82 55-60 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Begonia</i> sp. , <i>Bryopsis corticulans</i> , <i>Chlamydomonas reinhardtii</i> , psychrophilic <i>Chlamydomonas</i> sp. UWO241 and <i>Chlamydomonas</i> sp. ICE-MDV, <i>Chlorella vulgaris</i> , <i>Chromochloris zofingiensis</i> , <i>Colobanthus quitensis</i> Kunt Bartl, <i>Craterostigma pumilum</i> , <i>Cytisus cantabricus</i> (Wilk.) Rchb. F., <i>Dianthus caryophyllus</i> , <i>Drosera capensis</i> , <i>Euonymus japonicus</i> , <i>Fucus vesiculosus</i> , <i>Haematococcus pluvialis</i> , <i>Halomicronema hongdechloris</i> , <i>Hieracium pilosella</i> L., <i>Hordeum vulgare</i> , <i>Lasallia hispanica</i> , <i>Nicotiana tabacum</i> , <i>Oryza sativa</i> , <i>Pisum sativum</i> , <i>Marchantia polymorpha</i> (liverwort), <i>micro Nannochloropsis gaditana</i> , <i>Phaseolus vulgaris</i> , <i>Physcomitrella patens</i> , <i>Picea abies</i> , <i>Pinus strobus</i> , <i>Sinapsis alba</i> , <i>Spinacia oleracea</i> , <i>Synechococcus</i> PCC 7942, <i>Synechocystis</i> PCC 6803, <i>Syntrichia muralis</i> (Hedw.) Raab, <i>Scenedesmus obliquus</i> , <i>Ulva prolifera</i>
Predicted reactivity	Algae, <i>Bigeloviella natans</i> , <i>Cannabis sativa</i> , <i>Catalpa bungei</i> , <i>Citrus x limon</i> , Cyanobacteria, <i>Cyanidioschyzon merolae</i> strain 10D, <i>Lycopersicon esculentum</i> , <i>Panax ginseng</i> , <i>Picea spinulosa</i> , <i>Pinus thunbergii</i> , <i>Phaeodactylum tricornutum</i> , <i>Populus alba</i> , <i>Thermosynechococcus elongatus</i> (strain BP-1), <i>Triticum aestivum</i> Species of your interest not listed? Contact us
Not reactive in	<i>Chromera velia</i>
Additional information	Immunogold localization has been done in leaf material of <i>Arabidopsis thaliana</i> .

For high resolution images, please visit the specific product page at www.agrisera.com

Selected references

- [Kobayashi et al. \(2020\)](#). Relationship Between Glycerolipids and Photosynthetic Components During Recovery of Thylakoid Membranes From Nitrogen Starvation-Induced Attenuation in *Synechocystis* sp. PCC 6803. *Front Plant Sci.* 2020 Apr 15;11:432. doi: 10.3389/fpls.2020.00432. eCollection 2020.
- [Theurillat et al. \(2020\)](#). VIPP2 interacts with VIPP1 and HSP22E/F at chloroplast membranes and modulates a retrograde signal for HSP22E/F gene expression. *Plant Cell Environ.* 2020 Jan 29. doi: 10.1111/pce.13732.
- [Jokel et al. \(2020\)](#). Elimination of the flavodiiron electron sink facilitates long-term H₂ photoproduction in green algae. *Biotechnol Biofuels.* 2019 Dec 5;12:280. doi: 10.1186/s13068-019-1618-1.
- [Liu et al. \(2020\)](#). Acid treatment combined with high light leads to increased removal efficiency of *Ulva prolifera*. *Algal Research*, Volume 45, January 2020, 101745
- [Zhong et al. \(2019\)](#). Slower development of PSI activity limits photosynthesis during *Euonymus japonicus* leaf development. *Plant Physiol Biochem.* 2019 Mar;136:13-21. doi: 10.1016/j.plaphy.2019.01.004.
- [Roth et al. \(2019\)](#). Regulation of Oxygenic Photosynthesis during Trophic Transitions in the Green Alga *Chromochloris zofingiensis*. *Plant Cell.* 2019 Feb 20. pii: tpc.00742.2018. doi: 10.1105/tpc.18.00742.
- [Bastow et al. \(2018\)](#). Vacuolar Iron Stores Gated by NRAMP3 and NRAMP4 Are the Primary Source of Iron in Germinating Seeds. *Plant Physiol.* 2018 Jul;177(3):1267-1276. doi: 10.1104/pp.18.00478.
- [Kato et al. \(2018\)](#). Stepwise evolution of supercomplex formation with photosystem I is required for stabilization of chloroplast NADH dehydrogenase-like complex: Lhca5-dependent supercomplex formation in *Physcomitrella patens*. *Plant J.* 2018 Sep 3. doi: 10.1111/tj.14080.
- [Zhang et al. \(2018\)](#). VIRESCENT-ALBINO LEAF 1 regulates leaf colour development and cell division in rice. *J Exp Bot.* 2018 Aug 8. doi: 10.1093/jxb/ery250.
- [Giovanardi et al. \(2018\)](#). In pea stipules a functional photosynthetic electron flow occurs despite a reduced dynamics of LHCII association with photosystems. *Biochim Biophys Acta.* 2018 May 24. pii: S0005-2728(18)30129-4. doi: 10.1016/j.bbabi.2018.05.013.
- [Pao et al. \(2018\)](#). Lamelloplasts and minichloroplasts in Begoniaceae: iridescence and photosynthetic functioning. *J Plant Res.* 2018 Mar 2. doi: 10.1007/s10265-018-1020-2. (ImmunoGold)
- [He et al. \(2018\)](#). FRUCTOKINASE-LIKE PROTEIN 1 interacts with TRXz to regulate chloroplast development in rice. *J Integr Plant Biol.* 2018 Feb;60(2):94-111. doi: 10.1111/jipb.12631.
- [Myouga et al. \(2018\)](#). Stable accumulation of photosystem II requires ONE-HELIX PROTEIN1 (OHP1) of the light harvesting-like family. *Plant Physiol.* 2018 Feb 1. pii: pp.01782.2017. doi: 10.1104/pp.17.01782.
- [Muneer et al. \(2018\)](#). Proteomic Analysis Reveals the Dynamic Role of Silicon in Alleviation of Hyperhydricity in Carnation Grown In Vitro. *Int. J. Mol. Sci.* 2018, 19(1), 50; doi:10.3390/ijms19010050.
- [Schöttler et al. \(2017\)](#). The plastid-encoded Psal subunit stabilizes photosystem I during leaf senescence in tobacco. *J Exp Bot.* 2017 Feb 1;68(5):1137-1155. doi: 10.1093/jxb/erx009.
- [Fu et al. \(2017\)](#). Redesigning the QA binding site of Photosystem II allows reduction of exogenous quinones. *Nat Commun.* 2017 May 3;8:15274. doi: 10.1038/ncomms15274. (*Chlamydomonas reinhardtii*)
- [Sakuraba et al. \(2017\)](#). Rice Phytochrome-Interacting Factor-Like1 (OsPIL1) is involved in the promotion of chlorophyll biosynthesis through feed-forward regulatory loops. *Journal of Experimental Botany* doi:10.1093/jxb/erx231.
- [Gandini et al. \(2017\)](#). The transporter SynPAM71 is located in the plasma membrane and thylakoids, and mediates manganese tolerance in *Synechocystis* PCC6803. *New Phytol.* 2017 Mar 20. doi: 10.1111/nph.14526. (BN-PAGE)
- [Miguez et al. \(2017\)](#). Diversity of winter photoinhibitory responses: A case study in co-occurring lichens, mosses, herbs and woody plants from subalpine environments. *Physiol Plant.* 2017 Feb 14. doi: 10.1111/pl.12551.
- [Mazur et al. \(2016\)](#). Overlapping toxic effect of long term thallium exposure on white mustard (*Sinapis alba* L.) photosynthetic activity. *BMC Plant Biol.* 2016 Sep 2;16(1):191. doi: 10.1186/s12870-016-0883-4.
- [Yoshida et al. \(2016\)](#). Hisabori T1. Two distinct redox cascades cooperatively regulate chloroplast functions and sustain plant viability. *Proc Natl Acad Sci U S A.* 2016 Jul 5;113(27):E3967-76. doi: 10.1073/pnas.1604101113. Epub 2016 Jun 22.
- [Gerotto et al. \(2016\)](#). Flavodiiron proteins act as safety valve for electrons in *Physcomitrella patens*. *PNAS* DOI 10.1073.
- [Pavlovič et al. \(2016\)](#). A carnivorous sundew plant prefers protein over chitin as a source of nitrogen from its traps. *Plant Physiol Biochem.* 2016 Mar 5;104:11-16. doi: 10.1016/j.plaphy.2016.03.008
- [Pavlovič et al. \(2016\)](#). Light-induced gradual activation of photosystem II in dark-grown Norway spruce seedlings. *Biochim Biophys Acta.* 2016 Feb 18. pii: S0005-2728(16)30028-7. doi: 10.1016/j.bbabi.2016.02.009.

For high resolution images, please visit the specific product page at www.agrisera.com


Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Application example



2 µg of total protein from (1) *Arabidopsis thaliana* leaf, (2) *Hordeum vulgare* leaf, (3) *Chlamydomonas reinhardtii* total cell, (4) *Synechococcus* sp. 7942 total cell all extracted with Protein Extraction Buffer, PEB ([AS08 300](#)), were separated on **4-12%** NuPage (Invitrogen) **LDS-PAGE** and blotted 1h to **PVDF**. Blots were blocked immediately following transfer in 2% blocking reagent in 20 mM Tris, 137 mM sodium chloride pH 7.6 with 0.1% (v/v) Tween-20 (TBS-T) for 1h at room temperature with agitation. Blots were incubated in the primary antibody at a dilution of 1: 10 000 for 1h at room temperature with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at room temperature with agitation. Blots were incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, recommended secondary antibody [AS09 602](#)) diluted to 1:50 000 in 2% blocking solution for 1h at room temperature with agitation. The blots were washed as above and developed for 5 min with chemiluminescence detection reagent according to the manufacturer's instructions. Images of the blots were obtained using a CCD imager (FluorSMax, Bio-Rad) and Quantity One software (Bio-Rad). Exposure time was 10 seconds.