

# Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

Product no **AS07 257**

## PRK ribulose-5-P-kinase | Phosphoribulokinase

### Product information

<b>Immunogen</b>	KLH-conjugated peptide derived from known PRK sequences including <i>Arabidopsis thaliana</i> UniProt: <a href="#">P25697</a> , TAIR: <a href="#">At1g32060</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified serum in PBS, pH 7.4
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µg
<b>Reconstitution</b>	For reconstitution add 50 µl of sterile water.
<b>Storage</b>	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
<b>Additional information</b>	Antibody can be used as a marker of chloroplast stroma.

### Application information

<b>Recommended dilution</b>	1 : 1000 (WB)
<b>Expected   apparent MW</b>	44   39 kDa ( <i>A. thaliana</i> )
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Chlamydomonas reinhardtii</i> , <i>Hordeum vulgare</i> , <i>Synechocystis sp. PCC6803</i> , <i>Synechococcus PCC 7942</i> , <i>Thalassiosira pseudonana</i> , <i>Oryza sativa</i> , <i>Zea mays</i>
<b>Predicted reactivity</b>	<i>Glycine max</i> , <i>Hordeum vulgare</i> , <i>Lactuca sativa</i> , <i>Malus domestica</i> , <i>Micromonas sp.</i> , <i>Ostreococcus tauri</i> , <i>Physcomitrella patens</i> , <i>Populus trichocarpa</i> , <i>Spinacia oleracea</i> , <i>Solanum tuberosum</i> , <i>Sorghum bicolor</i> , <i>Synechocystis PCC 6803</i> , <i>Synechococcus elongatus</i> , <i>Zea mays</i> , <i>Vitis vinifera</i>  Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	Proteobacteria
<b>Additional information</b>	Antibody detects PRK using a load from 4-20 µg/well of a chloroplast fraction, incubation over night at 4 °C. For high resolution images, please visit the specific product page at <a href="http://www.agrisera.com">www.agrisera.com</a>
<b>Selected references</b>	<a href="#">Fukayama</a> et al. (2018). Expression level of Rubisco activase negatively correlates with Rubisco content in transgenic rice. <i>Photosynth Res.</i> 2018 May 30. doi: 10.1007/s11120-018-0525-9. <a href="#">Pérez-Ruiz</a> et al. (2017). NTRC-dependent redox balance of 2-Cys peroxiredoxins is needed for optimal function of the photosynthetic apparatus. <i>Proc Natl Acad Sci U S A.</i> 2017 Nov 7;114(45):12069-12074. doi: 10.1073/pnas.1706003114. <a href="#">Rai</a> et al. (2017). Real-time iTRAQ-based proteome profiling revealed the central metabolism involved in nitrogen starvation induced lipid accumulation in microalgae. <i>Sci Rep.</i> 2017 Apr 5;7:45732. doi: 10.1038/srep45732. (microalga, western blot) <a href="#">Nikkanen</a> et al. (2016). Crosstalk between chloroplast thioredoxin systems in regulation of photosynthesis. <i>Plant Cell Environ.</i> 2016 Aug;39(8):1691-705. doi: 10.1111/pce.12718.


# Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

## Applicaton example



WT-Col-0 *Arabidopsis thaliana* leaves were frozen in liquid nitrogen and soluble proteins were extracted in a buffer containing 50 mM HEPES, 5 mM NaCl and 10 mM MgCl<sub>2</sub>. 10 µg, 20 or 40 µg of eparated in SDS-PAGE (15% acrylamide with 6M urea) and blotted 1h to a PVDF membrane using Høefer semi-dry blotter. Blot was blocked with 4 % milk in TTBS for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1 000 overnight in +4 °C. The antibody solution was decanted and the blot was washed 3 x 5 min with TTBS at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera, [AS09 602](#)) diluted to 1:20 000 in 1% milk/TTBS for 2h at RT with agitation. The blot was washed 3x5 min in TTBS and 1x5 min in TBS at RT with agitation and developed for 5 min with chemiluminescent detection reagent, according to the manufacturer's instructions.

Courtesy of Lauri Nikkanen, University of Turku, Finland