

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS03 037-ALP**

RbcL | Rubisco large subunit, form I, ALP-conjugated (40 µg)

Product information

Immunogen	KLH-conjugated synthetic peptide conserved across all known plant, algal and (cyano)bacterial RbcL protein sequences (form I L8S8 and form II L2), including <i>Arabidopsis thaliana</i> AtCg00490 , <i>Hordeum vulgare</i> P05698 , <i>Oryza sativa</i> P0C510 , <i>Chlamydomonas reinhardtii</i> P00877 , <i>Synechococcus</i> PCC 7920 A5CKC5
Host	Rabbit
Clonality	Polyclonal
Purity	Affinity purified serum
Format	Liquid in PBS pH 7.4, conjugated to ALP
Quantity	40 µg
Storage	Store at 4°C for 12-18 months. A preservative may be added for long time storage up to 2 years.

Additional information	Anti-RbcL can be used as a cellular [compartment marker] of plastid stroma (cytoplasm in cyanobacteria) and detects RbcL protein from 31.25 fmoles. As both forms (I and II) are detected it is suitable for work with samples from Dinoflagellates, Haptophytes and Ochrophytes (diatoms, Raphidophytes, brown algae) as well as higher plants. This antibody together with Agrisera Rubisco protein standard is very suitable to quantify Rubisco in plant and algal samples.
-------------------------------	---

Application information

Recommended dilution	1 : 5000-10 000 (WB)
Expected apparent MW	52.7 kDa (<i>Arabidopsis thaliana</i>), 52.5 kDa (cyanobacteria), 52.3 (<i>Chlamydomonas reinhardtii</i>)
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Apium graveolens</i> , <i>Artemisia annua</i> , <i>Baculogypsina sphaerulata</i> (benthic foraminifer), <i>Bienertia sinuspersici</i> , <i>Cicer arietinum</i> , <i>Chlamydomonas raudensis</i> , <i>Chlamydomonas reinhardtii</i> , <i>Colobanthus quitensis</i> Kunt Bartl, <i>Cyanophora paradoxa</i> , <i>Cylindrospermopsis raciborskii</i> CS-505, <i>Emiliana huxleyi</i> , <i>Euglena gracilis</i> , <i>Fraxinus mandshurica</i> , <i>Fucus vesiculosus</i> , <i>Glycine max</i> , <i>Gonyaulax polyedra</i> , <i>Guzmania</i> hybrid, <i>Heterosigma akashiwo</i> , <i>Karenia brevis</i> (C.C.Davis) s) G.Hansen & Ø.Moestrup (Wilson isolate), <i>Liquidambar formosana</i> , <i>Micromonas pusilla</i> , <i>Nicotiana benthamiana</i> , <i>Physcomitrella patens</i> , <i>Porphyra</i> sp., <i>Schima superba</i> , <i>Stanleya pinnata</i> , <i>Spinacia oleracea</i> , lichens, <i>Symbiodinium</i> sp., <i>Synechococcus</i> PCC 7942, <i>Thalassiosira pseudonana</i> , <i>Thermosynechococcus elongatus</i> , <i>Prochlorococcus</i> sp. (surface and deep water ecotype), <i>Triticum aestivum</i> , dinoflagellate endosymbionts (genus <i>Symbiodinium</i>), extreme acidophilic verrucomicrobial methanotroph <i>Methylacidiphilum fumariolicum</i> strain SolV, <i>Thalassiosira punctigera</i> , <i>Vitis vinifera</i>
Predicted reactivity	Alpha-proteobacteria, Algae (brown and red), Dicots, Beta-proteobacteria, Conifers, Cryptomonads, Cyanobacteria (prochlorophytes), Gamma-proteobacteria, Liverworts, Monocots, Mosses, <i>Suaeda glauca</i> , <i>Welwitschia</i> ; <i>Nannochloropsis</i> sp. Species of your interest not listed? Contact us
Not reactive in	No confirmed exceptions from predicted reactivity are currently known.