

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS07 212**

VDAC1-5 | Voltage-dependent anion-selective channel protein 1-5

Product information

Background	VDAC proteins are porin-type, beta-barrel diffusion pores. Prominently localized in the outer mitochondrial membrane and involved in metabolite exchange.
Immunogen	KLH-conjugated peptide conserved in all known higher plant VDAC proteins including <i>Arabidopsis thaliana</i> VDAC1 UniProt: Q9SRH5 , TAIR: AT3G01280 , VDAC2 UniProt F4K3R8-1 , TAIR: AT5G67500 , VDAC3 UniProt: Q9SMX3-1 , TAIR: AT5G15090 , VDAC4 UniProt: Q9FKM2-1 , TAIR: AT5G57490 , VDAC5 UniProt: Q9M2W6-1 , TAIR: AT3G49920
Host	Rabbit
Clonality	Polyclonal
Purity	Affinity purified serum in PBS pH 7.4
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water.
Storage	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
Tested applications	Blue-native PAGE (2D BN/SDS-PAGE), Western blot (WB)
Related products	AS07 212-ALP Anti-VDAC1-5 Voltage-dependent anion-selective channel protein 1, ALP-conjugated (40 µg) AS07 212-HRP Anti-VDAC1-5 Voltage-dependent anion-selective channel protein 1, HRP-conjugated (40 µg) AS04 054 Anti-AOX1/2 rabbit antibody, marker of mitochondrial inner membrane AS06 203A Anti-Ildh rabbit antibody, marker of mitochondrial matrix collection of antibodies to other mitochondrial proteins
Additional information	Cellular [compartment marker] of mitochondrial outer membrane for western blot.

Application information

Recommended dilution	1 : 500 (IL), 1 : 5000, 2-30 µg protein/lane (WB)
Expected apparent MW	29 kDa (for <i>Arabidopsis thaliana</i>)
Confirmed reactivity	<i>Arabidopsis thaliana</i> , <i>Beta vulgaris</i> , <i>Brassica oleracea</i> var. botrytis, <i>Brassica rapa</i> subsp. <i>rapa</i> , <i>Citrus sinensis</i> , <i>Fortunella margarita</i> Swingle, <i>Oryza sativa</i> , <i>Papaver</i> sp. pollen tubes (IL), <i>Spinacia oleracea</i> , <i>Physcomitrella patens</i>
Predicted reactivity	<i>Arabidopsis alpina</i> , <i>Aundo donax</i> , <i>Brachypodium distachyon</i> , <i>Brassica campestris</i> , <i>Brassica napus</i> , <i>Brassica rapa</i> subsp. <i>pekinensis</i> , <i>Capsella rubella</i> , <i>Citrus clementina</i> , <i>Eutrema salsugineum</i> , <i>Glycine max</i> , <i>Glycine soja</i> , <i>Gossypium arboreum</i> , <i>Hoedum vulgare</i> var. <i>distichum</i> , <i>Jatropha curcas</i> , <i>Medicago truncatula</i> , <i>Mesembryanthemum crystallinum</i> , <i>Morus notabilis</i> , <i>Nicotiana tabacum</i> , <i>Phaseolus coccineus</i> , <i>Phaseolus vulgaris</i> , <i>Pisum sativum</i> , <i>Plantago major</i> , <i>Prunus persica</i> , <i>Ricinus communis</i> , <i>Solanum lycopersicum</i> , <i>Solanum tuberosum</i> , <i>Sorghum bicolor</i> , <i>Theobroma cacao</i> , <i>Triticum aestivum</i> , <i>Vitis vinifera</i> , <i>Zea mays</i> Species of your interest not listed? Contact us
Not reactive in	<i>Chlamydomonas reinhardtii</i> , <i>Glycine max</i> , <i>Zea mays</i> , diatoms, <i>Saccharomyces cerevisiae</i>

Additional information

Amount of mitochondrial fraction detected by anti-VDAC1 antibody was from 2-10 µg.

Immunolocalization method description and images are available [here](#)

Blue-native (2D BN/SDS-PAGE) methodology is described in Piechota et al. 2010


For high resolution images, please visit the specific product page at www.agrisera.com

Selected references

- [Tarasenko](#) et al. (2020). Plant mitochondrial subfractions have different ability to import DNA. *Theor. Exp. Plant Physiol.* doi.org/10.1007/s40626-020-00167-w
- [Garmash](#) et al. (2020). Altered levels of AOX1a expression result in changes in metabolic pathways in *Arabidopsis thaliana* plants acclimated to low dose rates of ultraviolet B radiation. *Plant Sci.* 2020 Feb;291:110332. doi: 10.1016/j.plantsci.2019.110332.
- [Bai](#) et al. (2019). Overexpression of soybean GmPLD enhances seed oil content and modulates fatty acid composition in transgenic *Arabidopsis*. *Plant Science* Volume 290, January 2020, 110298.
- [Klinger](#) et al. (2019). The signal distinguishing between targeting of outer membrane β -barrel protein to plastids and mitochondria in plants. *Biochim Biophys Acta Mol Cell Res.* 2019 Jan 8;1866(4):663-672. doi: 10.1016/j.bbamcr.2019.01.004.
- [Zhu](#) et al. (2018). A comprehensive proteomic analysis of elaioplasts from citrus fruits reveals insights into elaioplast biogenesis and function. *Hortic Res.* 2018 Feb 7;5:6. doi: 10.1038/s41438-017-0014-x.
- [Kang](#) et al. (2018). Autophagy-related (ATG) 11, ATG9 and the phosphatidylinositol 3-kinase control ATG2-mediated formation of autophagosomes in *Arabidopsis*. *Plant Cell Rep.* 2018 Jan 19. doi: 10.1007/s00299-018-2258-9.
- [Wang](#) and Auwerx (2017). Systems Phytohormone Responses to Mitochondrial Proteotoxic Stress. *Mol Cell.* 2017 Nov 2;68(3):540-551.e5. doi: 10.1016/j.molcel.2017.10.006.
- [Yin](#) et al. (2016). Comprehensive Mitochondrial Metabolic Shift during the Critical Node of Seed Ageing in Rice. *PLoS One.* 2016 Apr 28;11(4):e0148013. doi: 10.1371/journal.pone.0148013. eCollection 2016.
- [de Michele](#) et al. (2016). Free-Flow Electrophoresis of Plasma Membrane Vesicles Enriched by Two-Phase Partitioning Enhances the Quality of the Proteome from *Arabidopsis* Seedlings. *J Proteome Res.* 2016 Mar 4;15(3):900-13. doi: 10.1021/acs.jproteome.5b00876. Epub 2016 Feb 4.
- [Li](#) et al. (2015). A Chaperone Function of NO CATALASE ACTIVITY1 Is Required to Maintain Catalase Activity and for Multiple Stress Responses in *Arabidopsis*. *Plant Cell.* 2015 Feb 19. pii: tpc.114.135095.
- [Rurek](#) et al. (2015). Biogenesis of mitochondria in cauliflower (*Brassica oleracea* var. botrytis) curds subjected to temperature stress and recovery involves regulation of the complexome, respiratory chain activity, organellar translation and ultrastructure. *Biochim Biophys Acta.* 2015 Jan 21. pii: S0005-2728(15)00016-X. doi: 10.1016/j.bbabi.2015.01.005.
- [Armbruster](#) et al. (2014). Ion antiport accelerates photosynthetic acclimation in fluctuating light environments. *Nat Commun.* 2014 Nov 13;5:5439. doi: 10.1038/ncomms6439
- [Hsueh](#) et al. (2014). The chloroplast outer envelope protein P39 in *Arabidopsis thaliana* belongs to the Omp85 protein family. *Proteins.* 2014 Nov 17. doi: 10.1002/prot.24725.
- [Takahashi](#) et al. (2014). Transport of rice cyclobutane pyrimidine dimer (CPD) photolyase into mitochondria relies on a targeting sequence located in its C-terminal internal region.
- [Alcántar-Aguirre](#) et al. (2013). ATP produced by oxidative phosphorylation is channeled toward hexokinase bound to mitochondrial porin (VDAC) in beetroots (*Beta vulgaris*). *Planta*, March 17.

For high resolution images, please visit the specific product page at www.agrisera.com

Application example



Crude membrane proteins were separated on 12% SDS-PAGE and blotted 1h to PVDF. Blots were blocked immediately following transfer in 5% blocking reagent (BioRad, 170-6404) in 50 mM Tris, 150 mM sodium chloride pH 7.5 with 0.1% (v/v) Tween-20 (TBS-T) for 1h at room temperature with agitation. Blots were incubated in the primary antibody in 1: 5000 dilution for over-night at 4°C with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed 3 times for 5 min in TBS-T at room temperature with agitation. Blots were incubated in secondary antibody (Goat anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1:5000 in 0.2% blocking solution for 1h at room temperature with agitation. The blots were washed as above and developed for 1~2 min with chemiluminescent detection reagent, according the manufacturers instructions. Images of the blots were obtained using a CCD imager (LAS4000 GE) and by ImageQuant software (GE).

Arabidopsis thaliana membrane extraction and SDS-PAGE analysis About 200 mg (gFW) *Arabidopsis* seedlings (3-week-old), grown on 1% MS-agar plates, was ground with mortar and pestle in the presence of 2 ml extraction buffer [75 mM MOPS-KOH, 0.6 M Sucrose, 4 mM EDTA, 0.2% PVP-40, 0.2% BSA, 8 mM L-cystein, pH 7.6] and the protease inhibitor cocktail 'complete Mini' from Roche Diagnostics GmbH (Mannheim,

Agrisera


This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Germany). Crude membrane extracts were prepared essentially as described in Colas des Francs-Small et al. (2012). The membranous fraction was obtained by centrifugation at 22,000 g for 10 min at 4°C. The pellet containing the crude membranous fraction was washed twice with wash buffer [37.5 mM MOPS-KOH, 0.3 M Sucrose, 2 mM EDTA pH 7.6]. The samples were kept frozen at -80°C until used. For SDS-PAGE, an aliquot equivalent to 10 mg (i.e. 1x dilution) of crude Arabidopsis membrane extracts was solubilized in 3x Laemmli sample buffer (Bio-Rad) and the proteins were analyzed by SDS-gel electrophoresis

Courtesy of Dr. Oren Ostersetzer, The Hebrew University of Jerusalem, Israel



Fixation and Immunolocalization

(A) full confocal stacks; **(B)** Single confocal section

Pollen tubes were fixed in 400 μ M 3-maleimodobenzoic acid N-hydroxysuccinimide ester (MBS, Pierce) for 6 min at 20°C, followed by 2% formaldehyde (1 h, 4°C). Cells were washed three times in 1x TBS then once in MES buffer (15 mM MES, pH 5.0), then incubated in 0.05% cellulose/0.05% macerozyme with 0.1% Triton X-100 in MES buffer containing 0.1 mM PMSF and 1 % BSA for 15 min. Cells were washed once in MES, then twice in TBS and then incubated in blocking solution (1% BSA in TBS) for 30 min at room temperature. Pollen was incubated with anti-VDAC1 antibodies diluted in blocking solution (at 1:500) overnight at 4°C. Following TBS washes pollen was then incubated with the secondary antibody for 1.5 h at room temperature followed by further TBS washes. Pollen tubes were mounted on slides with 5 μ L of Vectashield + DAPI (Vector Laboratories, USA) and coverslips sealed with nail varnish.

Method taken from Poulter et al (submitted) Actin-binding proteins implicated in formation of the punctate actin foci stimulated by the self-incompatibility response in Papaver. Submitted to Plant Physiology.

Courtesy Professor Noni Franklin-Tong, University of Birmingham, UK