

# Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

Product no **AS06 167**

## PsbP | 23 kDa protein of the oxygen evolving complex (OEC) of PSII (anti-peptide)

### Product information

<b>Immunogen</b>	KLH-conjugated synthetic peptide derived from PsbP protein of <i>Arabidopsis thaliana</i> <a href="#">Q42029</a> , <a href="#">At1g06680</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	100 µl
<b>Reconstitution</b>	For reconstitution add 100 µl of sterile water.
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
<b>Additional information</b>	This product can be sold containing ProClin if requested.

### Application information

<b>Recommended dilution</b>	1 : 2000 (WB)
<b>Expected   apparent MW</b>	28   23 kDa ( <i>Arabidopsis thaliana</i> )
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Hordeum vulgare</i> , <i>Nicotiana tabacum</i> , <i>Picea abies</i> , <i>Pisum sativum</i> , <i>Physcomitrella patens</i>
<b>Predicted reactivity</b>	<i>Oryza sativa</i> , <i>Solanum tuberosum</i> , <i>Triticum aestivum</i> , <i>Picea sitchensis</i> , <i>Populus balsamifera</i> Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	<i>Chlamydomonas reinhardtii</i> , <i>Synechococcus</i> sp. PCC 7942, <i>Zostera marina</i>
<b>Additional information</b>	If you work with <i>Chlamydomonas reinhardtii</i> , please use following PsbP antibody: <a href="#">AS06 142-23</a> For high resolution images, please visit the specific product page at <a href="http://www.agrisera.com">www.agrisera.com</a>
<b>Selected references</b>	<a href="#">Pavlovič</a> et al. (2016). Light-induced gradual activation of photosystem II in dark-grown Norway spruce seedlings. <i>Biochim Biophys Acta</i> . 2016 Feb 18. pii: S0005-2728(16)30028-7. doi: 10.1016/j.bbabi.2016.02.009. <a href="#">Albanese</a> et al. (2016). Isolation of novel PSII-LHCII megacomplexes from pea plants characterized by a combination of proteomics and electron microscopy. <i>Photosynth Res</i> . 2016 Jan 9. <a href="#">Grassl</a> et al. (2012). Early events in plastid protein degradation in stay-green <i>Arabidopsis</i> reveal differential regulation beyond the retention of LHCII and chlorophyll. <i>J. Proteome Res</i> . October 2. <a href="#">Lang</a> et al. (2011). Simultaneous isolation of pure and intact chloroplasts and mitochondria from moss as the basis for sub-cellular proteomics. <i>Plant Cell Rep</i> . Feb;30(2):205-15. (reactivity confirmed for <i>Physcomitrella patens</i> ).


### Application example

# Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)



**2 µg of total protein** from (1) *Arabidopsis thaliana* leaf extracted with **Protein Extraction Buffer, PEB (AS08 300)**, (2) *Hordeum vulgare* leaf extracted with PEB, (3) *Chlamydomonas reinhardtii* total cell extracted with PEB, (4) *Synechococcus* sp. 7942 total cell extracted with PEB, were separated on **4-12% NuPage (Invitrogen) LDS-PAGE** and blotted 1h to **PVDF**. Blots were blocked immediately following transfer in 2% blocking reagent in 20 mM Tris, 137 mM sodium chloride pH 7.6 with 0.1% (v/v) Tween-20 (TBS-T) for 1h at room temperature with agitation. Blots were incubated in the primary antibody at a dilution of 1: 50 000 for 1h at room temperature with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at room temperature with agitation. Blots were incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1:10 000 in 2% blocking solution for 1h at room temperature with agitation. The blots were washed as above and developed for 5 min with chemiluminescent detection reagent according to the manufacturers instructions. Images of the blots were obtained using a CCD imager (FluorSMax, Bio-Rad) and Quantity One software (Bio-Rad).