


Product no **AS09 458S****PEPC | Phosphoenolpyruvate carboxylase positive control/quantitation standard****Product information****Format** | Lyophilized in glycerol.**Quantity** | 100 µl**Reconstitution** | For reconstitution, check the label on the tube. Please notice that this product contains 10% glycerol and might appear as liquid but is provided lyophilized.**Storage** | Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.**Additional information** | The PEPC protein standard can be used in a combination with Agrisera global PEPC antibody to quantitate PEPC from a wide range of species. [Global antibodies](#) are raised against highly conserved amino acid sequence.Quantitative western blot: [detailed method description](#), [video tutorial](#)**Application information****Recommended dilution** | Standard curve: 3 loads are recommended (0.5, 2 and 4 µl).
For most applications a sample load of 0.2 µg of chlorophyll/well will give a RbCL signal in this range.

Positive control: a 2 µl load per well is optimal for most chemiluminescent detection systems. Higher standard concentration needs to be used to allow detection by Coomassie stains. Such gels with higher standard concentration can not be used for quantitation using chemiluminescence.

This standard **is stabilized and ready** and does not require heating before loading on the gel.
Please note that this product contains 10% glycerol and might appear as liquid but is provided lyophilized. Allow the product several minutes to solubilize after adding water. Mix thoroughly but gently. Take extra care to mix thoroughly before each use, as the proteins tend to settle with the more dense layer after freezing.**Expected | apparent MW** | 110 | 105 kDa**Additional information** | **Concentration:** after re-constitution with sterile milliQ water final concentration of the standard is 0.15 pmoles/µl**Protein standard buffer composition:** Glycerol 10%, Tris Base 141 mM, Tris HCl 106 mM, LDS 2%, EDTA 0.51 mM, SERVA® Blue G250 0.22 mM, Phenol Red 0.175 mM, pH 8.5, 0.1 mg/ml PefaBloc protease inhibitor (Roche), 50 mM DTT.**This standard is ready-to-load and does not require any additions or heating.****This standard is stabilized and ready and does not require heating before loading on the gel.****Please note that this product contains 10% glycerol and might appear as liquid but is provided lyophilized.****Allow the product several minutes to solubilize after adding water. Mix thoroughly but gently. Take extra care to mix thoroughly before each use, as the proteins tend to settle with the more dense layer after freezing.****Application example**

5 µg total protein from 10-day old *Zea mays* leaves was loaded into Bolt 4-12% Bis-Tris Plus polyacrylamide gel (Invitrogen) in NuPAGE LDS sample buffer (1X, Invitrogen) and dithiothreitol (50 mM) along with four concentrations of PEPC standard (100, 200, 400 and 800 fmoles). The same concentrations of standard from lyophilized sample were also loaded into the same gel to evaluate the effectiveness of lyophilization process. Proteins were separated in Bolt MOPS SDS running buffer (1X, Invitrogen) at 200 V for 32 min using a Bolt Mini Gel Tank (Invitrogen). Proteins were then transferred onto 0.2-µm polyvinylidene fluoride membranes (PVDF, Immobilon) for 80 min at 20 V in NuPAGE transfer buffer (1X, Invitrogen) containing methanol (10%, v/v) and Bolt Antioxidant (Invitrogen) using Bolt Mini Blot Module (Invitrogen). Following the transfer, membrane was blocked for 1 h in 2% (w/v) membrane blocking agent dissolved in Tris-buffered saline solution containing Tween-20 (TBS-T; Tris, 20 mM; NaCl, 137 mM; Tween-20, 0.1% v/v). The membrane was treated with PEPC antibody (1:10,000, w/v in blocking solution, [AS09 458](#)) for

Agrisera

Part of Thermo Fisher Scientific

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

1h. Membrane was washed with TBS-T twice briefly, then once for 15 min and three times for 5 min each. Membrane was then incubated for 1 h with a horseradish peroxidase-conjugated goat anti-rabbit antibody (Agrisera; 1:25 000, w/v in blocking solution, [AS09_602](#)) followed by washing as described above. The signals were detected using ECL reagent and visualized using the Molecular Imager VersaDoc MP 4000 System (Bio-Rad).