

# Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

Product no **AS05 071**

## AtpH | ATP synthase subunit c (chloroplastic)

### Product information


<b>Immunogen</b>	Purified c subunit from <i>Spinacia oleracea</i> UniProt: <a href="#">P69447</a> .
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Serum
<b>Format</b>	Lyophilized
<b>Quantity</b>	200 µl
<b>Reconstitution</b>	For reconstitution add 200 µl of sterile water.
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.

**Additional information** | This product can be sold containing proclin if requested.

### Application information

<b>Recommended dilution</b>	1 : 1000-1 : 10 000 (WB)
<b>Expected   apparent MW</b>	8 kDa (for <i>Arabidopsis thaliana</i> )
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Nicotiana benthamina</i> , <i>Spinacia oleracea</i> , <i>Thermosynechococcus elongatus</i>
<b>Predicted reactivity</b>	Algae, <i>Cannabis sativa</i> , <i>Glycine max</i> , <i>Oryza sativa</i> , <i>Physcomitrella patens</i> , <i>Pisum sativum</i> , <i>Populus alba</i> , <i>Pinus thunbergii</i> , <i>Zea mays</i> , <i>Vitis vinifera</i>  Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known.
<b>Additional information</b>	Note that increased incubation at 95°C (20-30 min) prior to loading is recommended to break the multimeric c-mer structure, detection of partial ring structures (e.g. 5 or 6 subunits) may occur.  For high resolution images, please visit the specific product page at <a href="http://www.agrisera.com">www.agrisera.com</a>
<b>Selected references</b>	<a href="#">Nath et al. (2016)</a> . A Nitrogen-Fixing Subunit Essential for Accumulating 4Fe-4S-Containing Photosystem I Core Proteins. <i>Plant Physiol.</i> 2016 Dec;172(4):2459-2470. Epub 2016 Oct 26. <a href="#">Lawrence et al. (2010)</a> . Recombinant production and purification of the subunit c of chloroplast ATP synthase. <i>Protein Expression and Purification</i> 76: 15-24.

### Application example



# Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

0.7 µg of purified ATP-synthase complex from *Spinacia oleracea* (CF0F1) (1), 15.3 µg of purified ATP-synthase complex (CF0F1) from *Nicotiana benthamiana* (2) and 48.6 µg of *Thermosynechococcus elongatus* thylakoid preparation (3) were separated on 12% polyacrylamide gel and blotted on PVDF membrane. Filters were blocked (0.5h), incubated with 1: 1000 anti-AtpH antibodies (1h), followed by incubation with 1: 5 000 secondary anti-rabbit antibody (1.25h), coupled to HRP and visualized with chemiluminescence detection reagent.