


Product no **AS10 652****Anti-Cu/ZnSOD | Cu/Zn superoxide dismutase****Product information**

Immunogen	15 amino acids synthetic peptide designed from olive pollen cytosolic Cu/Zn cDNA consensus sequences (GenBank Accessions EU250757 to EU250797), UniProt Q8L5E0
Host	Chicken
Clonality	Polyclonal
Purity	Purified, total IgY in 10 mM Tris, 150 mM NaCl, 0.02% sodium azide.
Format	Liquid
Quantity	100 µl
Storage	Store at 4°C. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
Additional information	Total IgY concentration is 2.3 mg/ml. Reaction of the antibody to chloroplastic SOD isoform has not been determined yet.

Application information

Recommended dilution	1 : 1500 with Alexa 488 (WB)
Expected apparent MW	15.3 16 kDa (<i>Olea europaea</i> L.)
Confirmed reactivity	<i>Arabidopsis thaliana</i> (leaves), <i>Lilium longiflorum</i> , <i>Olea europaea</i> (pollen), <i>Oryza sativa</i> , <i>Spinacia oleracea</i> (leaves), <i>Solanum lycopersicum</i> , <i>Zea mays</i>
Predicted reactivity	<i>Ananas ananas</i> , <i>Betula pendula</i> , <i>Camellia sinensis</i> , <i>Codonis lanceolata</i> , <i>Cucurbita ficifolia</i> , <i>Helianthus sp.</i> , <i>Hordeum vulgare</i> , <i>Lycopersicum esculentum</i> , <i>Plantago major</i> , <i>Populus trichocarpa</i> , <i>Solanum nigrum</i> , <i>Solanum tuberosum</i> , <i>Solidago sp.</i> , <i>Vitis vinifera</i> Species of your interest not listed? Contact us
Not reactive in	<i>Helianthus annuus</i> , <i>Nicotiana tabacum</i> , <i>Marchantia polymorpha</i> , <i>Musa acuminata</i> , <i>Vicia faba</i>
Additional information	Note: Antibody recognizes two to three isoforms of Cu/Zn SOD in olive pollen depending on the olive cultivar
Selected references	Adhikari et al. (2018). Sulfate improves cadmium tolerance by limiting cadmium accumulation, modulation of sulfur metabolism and antioxidant defense system in maize. <i>Environmental and Experimental Botany</i> Volume 153, September 2018, Pages 143-162. Bastow et al. (2018). Vacuolar Iron Stores Gated by NRAMP3 and NRAMP4 Are the Primary Source of Iron in Germinating Seeds. <i>Plant Physiol.</i> 2018 Jul;177(3):1267-1276. doi: 10.1104/pp.18.00478. Alché et al. (1998). Identification and immunolocalization of superoxide dismutase isoenzymes of olive pollen". <i>Physiol. Plantarum</i> 104, 772-776.

Application example

25 µg of total protein from *Olea europaea* pollen were separated on 12% acrylamide gels and blotted onto PVDF. Membranes were blocked 1h with 2% low-fat milk powder in TBS-T (0.1% TWEEN 20) and probed with anti-Cu/Zn SOD (AS10 652, 1:1500, overnight, 4° C) and secondary antibody anti-chicken IgG conjugated to Alexa 488 (1:2000, 1 h) (Molecular Probes, Invitrogen). Signal was detected with a PharosSX fluorescence scanner (Bio-Rad).

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

The antibody recognizes a protein of c.a. 16 kDa. and another band likely corresponding to the protein dimer.

Courtesy Dr. Juan de Dios Alche, CSIC, Spain