

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS10 817**

Goat anti-Human IgG Fc, F(ab)² fragment, HRP conjugated, min. cross-reactivity bovine/mouse/rabbit serum

Product information

Immunogen | Purified human IgG

Host | Goat

Clonality | Polyclonal

Purity | Affinity purified goat IgG

Format | Lyophilized

Quantity | 0.5 mg

Reconstitution | For reconstitution add 0.55 ml of sterile water

Storage | Store lyophilized material at 2-8 °C.
For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20 °C, to prevent loss of enzymatic activity.
For example, if you have reconstituted 0.5 mg of antibody in 0.55 ml of sterile water add 0.55 ml of glycerol. Such solution will not freeze in -20 °C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.

Additional information | HRP-conjugate is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free
0.1 % (v/v) of Kathon CG is used as preservative. Use of sodium azide will inhibit enzyme activity of horseradish peroxidase

Application information

Recommended dilution | The optimal working dilution should be determined by the investigator.

Confirmed reactivity | Human IgG Fc (two heavy chains with constant domains)

Predicted reactivity | Human IgG Fc (two Heavy chains with constant domains)

Not reactive in | No confirmed exceptions from predicted reactivity are currently known.

Additional information | This antibody reacts with the heavy chains on human IgG based on immunoelectrophoresis.
No reactivity is observed with the light chains on human immunoglobulins or non-immunoglobulin human serum proteins or bovine, mouse or rabbit serum proteins or IgG F(ab)² fragment based on immunoelectrophoresis.
The antibody has been absorbed against solid phase normal bovine, mouse and rabbit serum in addition to mouse and rabbit IgG.