

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

product **AS16 3208**

TGA1 | TGACG motif-binding factor 1, bZIP transcription factor

product information

Background	TGA1 (TGACG motif-binding factor 1) is a transcriptional activator which binds to the 5'-TGACG-3' DNA sequence. Its binding to the as-1-like cis elements mediate salicylic acid- and auxin-inducible transcription. TGA1 interacts with NPR1 and is involved in the induction of the systemic acquired resistance (SAR).
Immunogen	KLH-conjugated peptide derived from protein sequence of <i>Arabidopsis thaliana</i> TGA1, UniProt: Q39237 , TAIR: At5g65210
Host	Rabbit
Clonality	Polyclonal
Purity	Affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water.
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.
Tested applications	Immunoprecipitation (IP), Western Blot (WB)
Related products	AS16 3209 Anti-TGA2 TGACG motif-binding factor 2, bZIP transcription factor, rabbit antibodies Antibodies to proteins involved in regulation of transcription Plant protein extraction buffer Secondary antibodies

Application information

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	42.1 kD
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Brassica sp.</i> , <i>Camelina sativa</i> , <i>Eutrema salsugineum</i> , <i>Gossypium arboreum</i> , <i>Gossypium raimondii</i> , <i>Jatropha curcas</i> , <i>Morus notabilis</i> , <i>Nelumbo nucifera</i> , <i>Populus sp.</i> , <i>Ricinus communis</i> , <i>Tarenaya hassleriana</i> , <i>Theobroma cacao</i> , <i>Vitis vinifera</i>
Additional information	TGA1 level in leaf may be too low for detection. Antibody is recognizing transiently expressed TGA1 with GFP tag.
Selected references	To be added when available, antibody released in January 2018.


Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Application example



50 µg of total protein from *Arabidopsis thaliana* seedlings of col-0, *tga1-1* and *Nicotiana tabacum* leaf transiently expressing TGA1-GFP (described sample order from left to right), extracted with extraction buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 5 mM EDTA, 0.1% Triton X-100, 0.2% Nonidet P-4) and denatured at 70 °C for 15 min were separated on 4-12% SDS-PAGE and blotted 1h to nitrocellulose membrane using tank transfer. Blots were blocked with 5% milk in TBS-T for 1h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1 000 at 4 °C overnight with agitation in TBS-T. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1:10 000 in for 1h at RT with agitation. The blot was washed as above and developed for 5 min with SuperSignal™ West Pico PLUS Chemiluminescent Substrate. Exposure time was 2min.

Courtesy of Dr. Jian Chen, Graduate Science Research Center Columbia, South Carolina, USA