

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS21 4570

Anti-APP-C31 | Caspase 3 cleaved APP, C terminal fragment

Product information

Immunogen Synthetic peptide corresponding to the N-terminal of human caspase 3-generated APP C-terminal 31 amino acids (aa 665-670) UniProt: P05067

Host Rabbit

Purity Serum. Contains 0.05% sodium azide.

Format Liquid

Quantity 100 μl

Reconstitution For reconstitution, add μl of sterile water

Storage Store at -20°C; make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution 1::500 - 1: 1000 (IL), 1:1000-1:3000 (WB)

Expected | apparent 86.9 | 100 kDa


Confirmed reactivity Human, Mouse, Rat

Predicted reactivity Species of your interest not listed? Contact us

Species of your interest flot listed: Outlact us

Selected references

Nishimura et al. (2002) Cell death induced by a caspase-cleaved transmembrane fragment of the Alzheimer amyloid precursor protein. Cell Death Differ. 2002 Feb;9(2):199-208. doi: 10.1038/sj.cdd.4400931. PMID: 11840170.



Sample: NT2 neurons

Fixation: 72 h after infection with adenovirus vector expressing Amyloid Precursor Protein Anti-APP-C31 antibody was used in dilution 1: 1000. Left image: stained for the N-terminus of APP with P2-1, mouse monoclonal antibody specific for N-terminus of APP.

Center image: stained for chromosomal DNA (Hoechst). Right image: stained for C-terminus of the caspase 3 cleaved APP with anti-APP-C31 antibody (ACT1).

Most of APP-accumulating neurons with shrunken and fragmented nuclei contain ACT1-immunoreactivity (arrows), but non-neuronal cells are hardly labeled with ACT1 (arrowheads).

The tissue was fixed with 4 % formaldehyde and permalized with acetone at -20° C, followed by blocking with 4 % non-fat milk and 5 % goat normal serum at 4 ° C/ON. Primart antibody was incubated at 1: 1000 1h/RT.